



PASPCR Newsletter

Volume 4 Number 2

June, 1996

Introduction . . .

by the Publications Committee

The **PASPCR Newsletter** is published quarterly and is intended to serve as a means of communication for the members of our Society. As such, we invite our membership to actively contribute to the *Newsletter*; help us to update the Job Listings, Calendar of Events, Meeting Reports, Abstracts in press and other items of general membership interest. If you attend a scientific meeting at which you heard about work which you think will be of interest to the membership of the **PASPCR**, please write a few paragraphs summarizing what was presented and share it with us. If you should have a change of affiliation or address, we'd like to know that, too. This is **your Newsletter**, and we depend upon you to help us make sure it best serves the Society's needs. Contributions and comments can be sent to any of the members of the Publications Committee.

NEW! WorldWideWeb Pages for the PASPCR. The PASPCR now has its own **WWW** home page. We plan this to be a major source of current information for the PASPCR membership. The address for the page is: <http://lenti.med.umn.edu/paspcr>. This site contains information on the goals of the society, future meetings, council information, past issues of the PASPCR newsletter as well as links to other sites including the InterPig DataBase, the International Pigment Cell Conference in Anaheim and the International Federation of Pigment Cell Societies (IFPCS).

We have now included the membership directory on that page; please notify us if you wish any or all of your information to be deleted or modified on that site.

The PASPCR WWW page system is still under construction and we want to know if there is any other information you would like located on this site.

Please check out the PASPCR web site and send any comments and/or suggestions to either the PASPCR WebMaster Bill Oetting at bill@lenti.med.umn.edu or to Vince Hearing at hearingv@dc37a.nci.nih.gov.

IN THIS ISSUE

Introduction	p 1
PASPCR Contact Information	p 2
Calendar of Events	p 2
Welcome to New Members	p 3
Corporate Sponsors	p 3
1996 XVI th Intl Pigment Cell Conf	p 3
Positions Wanted / Available	p 5
INTERPIG DataBase	p 5
Meeting Report - Intl Symposium on Melanogenesis and Melanoma	p 9
Bibliography	p 9

**PanAmerican Society for
Pigment Cell Research**

c/o **Dr. James J. Nordlund**
Department of Dermatology
University of Cincinnati
231 Bethesda Avenue
Cincinnati, OH 45267-0592
FAX: (513) 558-0198

Officers

Sally Frost-Mason
President
Richard A. King
President-Elect
James J. Nordlund
Secretary/Treasurer

Council Members

Raymond E. Boissy
Alan N. Houghton
Gert M. Jacobsohn
Kenneth A. Mason
Frank L. Meyskens, Jr.
David A. Norris
William S. Oetting
Walter C. Quevedo, Jr.
DeWayne Townsend

IFPCS Representative

Vincent J. Hearing
Past-President

The **PASPCR Newsletter** is published quarterly; for further information and/or to submit articles, contact the:

Publications Committee:

Dr. Kenneth A. Mason (chair)

University of Kansas
Department of Biochemistry
Lawrence, KS 66045
Phone: 913/864-4279
FAX: 913/864-5321

Frank L. Meyskens, Jr.

University of California - Cancer Center
101 City Drive
Orange, CA 92668
Phone: 714/456-6310
FAX: 714/456-5039

David A. Norris

Department of Dermatology
University of Colorado Medical Center
4200 East 9th Avenue
Denver, CO 80262
Phone: 303/372-1142
FAX: 303/372-1159

Calendar of Events :

June 2 - 6, 1996 Annual Meeting of the American Society for Biochemistry and Molecular Biology and the American Association of Immunologists, to be held in New Orleans, LA (contact: FASEB Office, 9650 Rockville Pike, Bethesda, MD 20814-3998; FAX: 301/530-7014)

June 15 - 16, 1996 5th Melanoma Workshop, to be held in Hamburg, Germany (contact: Holger Voigt, Melanoma Research Project, 31 B Dammtorstrasse, Hamburg D-20354, Germany; FAX: +49 40/348-0525)

Oct 29- Nov 3, 1996 XVIth International Pigment Cell Conference, to be held in Anaheim, California, (contact: MMC/UCI Center for Health Education, PO Box 1428, Long Beach, CA 90801-1428, FAX: 310/933-2012)

Dec 7 - 11, 1996 36th Annual Meeting of the American Society for Cell Biology and 6th International Congress on Cell Biology, to be held in San Francisco, CA, (contact: ASCB Secretariat, 9650 Rockville Pike, Bethesda, MD 20814-3992; FAX: 301/530-7139)

Jun 10 - 14, 1997 4th World Conference on Melanoma to be held in Sydney, Australia (contact: The Melanoma Foundation, PO Box M123, Camperdown, NSW 2050 Australia; FAX: +61 2/550-6316)

Jun 15- 18, 1997 VIIth PASPCR Annual Meeting, to be held in Providence, RI (contact: Dr. Walter C Quevedo, Jr., Brown University, Division of Biology and Medicine, Providence, RI 02912; FAX: 401/863-1971)

Jun 22 - 24, 1997 International Meeting "Pigmentary Disorders from a Global Perspectives" to be held in Bali, Indonesia (contact: Bureau PAOG, Tafelbergweg 25, 1105 BC Amsterdam, The Netherlands; FAX: +31 20/696-3229)

Welcome to New Members

by James J Nordlund

We welcome the following new member to the PASPCR . . .

Dong Fang	Mei-Yu Hsu	Marcus D. Johansen
Luix Eduardo M. Nery	Tomasz W. Panz	Brinda K. Rana
Deepa Rungta	Fan Yang	

If anyone is interested in joining our Society or wishes to sponsor a member, application forms can be obtained from Dr. James J. Nordlund at the PASPCR Secretary/Treasurer's office.

Corporate Sponsors

by James J Nordlund

The PASPCR would like to acknowledge and thank our Corporate Sponsors; the list below reflects contributions over the past 2 years. Financial gifts from these sponsors have allowed our Society to increase benefits to the membership far out of proportion to the actual dues collected from members. Monies contributed by these sponsors have been used over the years to support various PASPCR functions including our Young Investigator Award program, meeting travel stipends, annual meeting expenses and this Newsletter.

GOLD Corporate Patrons

ICN Pharmaceuticals, Inc
Lawrence M Gelb Research
Foundation of Clairol, Inc
Ortho Pharmaceutical Corp

SILVER Corporate Patrons

Avon Products, Inc
Chesebrough-Pond's USA Co
Procter and Gamble Co
Shiseido Co, Ltd

Corporate Patrons

Galderma Laboratories, Inc
Stiefel Laboratories

XVIth IPCC (International Pigment Cell Conference)

by Roger Bowers, Frank Meyskens

The XVIth International Pigment Cell Conference will be held from October 29th to November 3rd, 1996 at the Disneyland Hotel in Anaheim, California. Frank Meyskens is the Organizer of this meeting with Roger Bowers and Alistair Cochran serving as co-chairs of the Organizing Committee. The PASPCR has established a Web page that contains relevant information for this meeting; take a look at: "<http://lenti.med.umn.edu/paspcr/ipcc.htm>".

Authors of abstracts will be notified of their place in the program by mid-July. All keynote and invited platform speakers will be asked to submit a manuscript for publication in *Pigment Cell Research* in the upcoming year.

XVIth International Pigment Cell Conference - Tentative Conference Agenda

Tuesday, October 29, 1996

3:00 - 7:00 pm Pre-registration/View Exhibits
7:00 - 10:00 pm Welcome Reception: Fashion Show: "*Safe and Sexy in the Sun*"

Wednesday, October 30, 1996 Conference Attendees

7:00 - 8:00 am Registration/Continental Breakfast/View Exhibits
8:00 - 8:05 am Welcome: Chairman, Frank L. Meyskens, Jr.
Introduction: Laurel Wilkening, Chancellor, Univ of California, Irvine
8:05 - 8:35 am *Special Lecture*, R. Sherwood Rowland, Nobel Laureate, 1995, Chemistry
"Ozone Depletion, Ultraviolet Light, and the Pigment Cell"

Symposium I: Economic and Societal Implications of Melanin and Melanogenesis

8:35 - 9:00 am Keynote Speaker
9:00 - 10:30 am Invited and Competitive Abstract Speakers
10:30 - 11:00 am Break

11:00 - 12:30 pm **Workshop A: "Extracutaneous Melanin"**
 Posters and Discussion #1 TBN* (11:00 - 12:00 Viewing; 12:00 - 12:30 Discussion)
 12:30 - 2:00 pm Lunch on your own
Symposium II: Molecular Biology of Pigment Cells
 2:00 - 2:30 pm Keynote Speaker
 2:30 - 4:00 pm Invited and Competitive Abstract Speakers
 4:00 - 4:05 pm IFPCS InterPig Database on the WorldWideWeb: Vincent Hearing
 4:05 - 4:15 pm Break
 4:15 - 6:15 pm **Workshop B: "Regulating Mechanisms of Melanocyte Proliferation"**
 4:15 - 6:00 pm Posters and Discussion #2 TBN* (4:15 - 5:30 Viewing; 5:30 - 6:00 Discussion)
 5:30 - 7:00 pm **Workshop C: "Biophysics of Melanin"**
 6:15 pm Adjourn Free evening

Accompanying Guests (8:00 - 11:00 am) Welcome/Introduction: Buffet Breakfast;
 (12:00 - 5:00 pm) Group Activity

Thursday, October 31, 1996

7:00 - 8:00 am Continental Breakfast/View Exhibits
 8:00 - 8:30 am **Seiji Lectureship:** Introduction: Giuseppe Prota, President, IFPCS

Symposium III: Melanoma Research: Basic and Applied

8:30 - 9:00 am Keynote Speaker
 9:00 - 10:30 am Invited and Competitive Abstract Speakers
 10:30 - 11:00 am Break
 11:00 - 12:30 pm **Workshop D: "Control of Melanogenesis"**
 11:00 - 12:30 pm Simultaneous Business Meetings of Regional Societies
 12:30 - 2:00 pm Lunch on your own

Symposium IV: Photobiology of Melanocytes: Etiology and Prevention

2:00 - 2:30 pm Keynote Speaker
 2:30 - 4:00 pm Invited and Competitive Abstract Speakers
 4:00 - 7:00 pm **Workshop E: The "Blues" Symposium**
 4:00 - 7:00 pm Poster Viewing
 Adjourn - Free evening

Friday, November 1, 1996

7:00 - 8:00 am Continental Breakfast/View Exhibits
 8:00 - 8:30 am Introduction: Sally Frost-Mason, *President*, PASPCR
Gelb Lectureship: Seth Orlow

Symposium V: Melanogenesis and Pigmentary Disorders

8:30 - 9:00 am Keynote Speaker
 9:00 - 10:30 am Invited and Competitive Abstract Speakers
 10:30 - 11:00 am Break
 11:00 - 12:30 pm **Workshop F: "Biology and Biochemistry of Melanosomes"**
 11:00 - 12:30 pm Posters and Discussion #3 TBN* (11:00 - 12:00 Viewing; 12:00 - 12:30 Discussion)
 12:30 - 1:15 pm **Controversy Session:** "Semiquinone Radicals are not Important during Melanin Synthesis"
 Pro: Patrick Riley Con: Stan Pavel
 1:15 pm Adjourn Scientific Session
 1:15 - 6:30 pm Break
 6:30 - 7:30 pm Reception
 7:30 - midnight Banquet, Awards and Dancing

Saturday, November 2, 1996

7:00 - 8:00 am Continental Breakfast/View Exhibits
 8:00 - 8:30 am **Presidential Address:** Giuseppe Prota, President IFPCS

Symposium VI: Comparative Developmental Biology of Pigment Cells

8:30 - 9:00 am Keynote Speaker
 9:00 - 10:30 am Invited and Competitive Abstract Speakers
 10:30 - 11:00 am Break
 11:00 - 12:30 pm **Workshop G: "Genetic Aspects of Albinism"**
 11:00 - 12:30 pm Posters and Discussion #4 TBN* (11:00 - 12:00 Viewing; 12:00 - 12:30 Discussion)
 12:30 - 2:00 pm Lunch on your own
 2:00 - 4:00 pm **Educational Forum: "Living with the Sun"**
 4:00 - 6:00 pm Family Farewell Reception and Wine Tasting

Sunday, November 3, 1996

8:00 - 5:00 pm

1. *Satellite Meeting (all day): Classification of Cutaneous Melanoma:* Alistair Cochran
2. *Satellite Meeting (3 hours): Safety of Sunscreens and Tanning Parlors:* J.P. Césarini, et al.
3. *Satellite Meeting (3 hours): Ocular Melanin:* Giuseppe Prota

Workshops and poster and poster discussion sessions will be simultaneous. The poster sessions and discussions will feature areas that do not overlap with the workshop. The chairs of these sessions will be selected from submitted competitive abstracts and the Chairman in turn will organize this session with help from the Organizing Committee.

Positions - Wanted and Available :

Assistant Professor - The Department of Physiology at the University of Sao Paulo has an open position for assistant professor. Applications will be received until June 1st. A Ph.D. is required and post-doctoral experience is desirable. The selection will be held during the first week of July with the candidates' presence. The curriculum vitae including copies of the publications, the Ph.D. certificate and a research project on Human or Animal Physiology (including biochemical and biomolecular approaches) should accompany the application. A lecture (undergraduate level) on a picked out subject of one of three areas (Neurosensorial and Muscle Physiology; Cardiorespiratory and Excretory Physiology; Nutrition, Metabolism and Endocrine Physiology) chosen by the candidate should be given to the Selection Committee, that will also interview the candidates. Please contact: Ana Maria de L. Castrucci, Dep. Physiology, Inst. Biosciences, University of Sao Paulo, CP11176, CEP 05422-970, Sao Paulo, Brasil; Fax 5511 818-7422 phone 5511 818-7610 Email AMDLCAST@USP.BR

Predoctoral and Postdoctoral Positions - available for molecular biologists in the areas of drug discovery and metabolism research. Requires experience in gene cloning, DNA sequencing, recombinant protein expression and cell culture methods. Prior experience in dermatology research is desirable. Southern Research Institute is a diversified research and development organization. Our Life Sciences Division provides comprehensive preclinical drug development and testing capabilities as well as basic research in drug design and synthesis, pharmaceutical formulations, toxicology, virology, microbiology, and pharmacology. To apply, send resume or curriculum vitae to: Southern Research Institute, Attention: Suzann Allen, Human Resources, Department 118, P.O. Box 55305, Birmingham, AL, 35255-5305.

Faculty Position - Massachusetts General Hospital, Harvard Medical School, Cutaneous Biology Research Center. The Cutaneous Biology Research Center (CBRC) seeks a molecular, cellular or developmental biologist to establish a program in fundamental research relevant to skin pigmentation. Areas of research can include but are not limited to pigment synthesis and transfer in melanocytes, genetics of mouse coat color and development/migration of neural crest cells. Applicants must have a Ph.D. and/or M.D. degree and relevant postdoctoral experience. Only applicants with a strong research record and the potential to develop extramurally supported research programs will be considered. Individuals with a demonstrated ability to develop imaginative approaches to important biological questions are particularly encouraged to apply. Rank/salary/start-up funds and space are negotiable depending on experience and qualifications. The CBRC occupies 45,000 square feet of fully equipped laboratory space in a new multidisciplinary research facility. Interested individuals should send curriculum vitae, reprints, a statement of research and future directions, along with the names, addresses and telephone numbers of three references to: Dr. Paul F. Goetinck, Chair, Faculty Search Committee, Cutaneous Biology Research Center, Massachusetts General Hospital - East, Building 149, 13th Street, Charlestown, MA 02129

INTERPIG DataBase

by Vincent Hearing

The INTERPIG database is on the InterNet! You can now access the InterPig DataBase at the following address: <http://lenti.med.umn.edu/paspcr/interpig.html>. Please note that as of this time, I estimate that less than 5% of the various IFPCS members have contributed entries. Think of how useful and complete this list would be if everyone took the time to supply their own information. Please take a moment to fill out the database data entry form (either online through the Web page or

via Email) and send it back to Dr. Hearing. Please contact Vince Hearing or Bill Oetting if you need more information about these mechanisms of submission.

Meeting Report

by Kazunori Urabe

International Symposium on "Melanogenesis and Malignant Melanoma: Biochemistry, Cell Biology, Molecular Biology, Pathophysiology, Diagnosis and Treatment". - Fukuoka, Japan December 5-6, 1995

This symposium was held by Yoshiaki Hori, M.D., Professor and Chairman, Department of Dermatology, Faculty of Medicine, Kyushu University, to discuss new findings about melanogenesis, biochemistry of melanin, genetics of pigmented disorders, immunology of pigmented disorders and malignant melanoma, clinical features of malignant melanoma, prevention of malignant melanoma and treatment of malignant melanoma. Sixteen investigators were invited from abroad to report their new data on their fields and thirty Japanese ones were also invited to attend the meeting and sixteen of them presented their findings. The symposium was opened in the international conference room in the new modern building which was built for the international congresses a half year ago. Exact respected schedule by the speakers gave opportunity to everyone to hear and discuss what they were interested.

There were five sessions in this symposium. The first session dealt with the biochemistry and molecular biology in benign and malignant pigment tissues. In this session, the relationship between genetic alterations in melanogenic proteins and the clinical pigment disorders were reported. And the biological characters of melanogenic cells affected by surrounding cells and the environmental stimulants were investigated. In the second session, immunological approaches for malignant melanoma were presented. Several specific antigens in melanoma were reported and the possibility for clinical application were discussed. In the third session, the characteristic nature of melanoma cell were analyzed intensively. In the fourth session, some of new techniques for the diagnosis of malignant melanoma were presented. And the last session included the several new therapeutic methods; biochemotherapy, isolated limb perfusion, hyperthermia, and new products targeting melanoma. All papers were summarized here. A full listing of abstract is available on request. However, papers reported in this meeting will be published in a book from Elsevier Science soon.

1) Biochemistry and Molecular Biology of Melanogenesis in Benign and Malignant Pigment Tissues

Matsunaga J from Akita University of Medicine reported the mutations in the tyrosinase gene causing tyrosinase-negative oculocutaneous albinism in Japanese. In these nine patients, mutations of the tyrosinase gene were found at codon 77, codon 278, codon 310 and codon 431 of the gene. He suggested that in these mutations, codon 77 and codon 310 might be major mutation sites in Japanese patients. King RA from University of Minnesota demonstrated eleven mutations (9 unique and 2 previous reported) of the P gene which is responsible for tyrosinase positive oculocutaneous albinism. Hermansky-Pudlak syndrome was also intensively studied by his group, which is a type of albinism associated with storage-pool deficient platelets and with the production of ceroid. Linkage analysis of the family revealed that the gene was located on chromosome 10q. Further studies were going on to clone the gene. I presented the data about the regulation of *microphthalmia (mi)* gene of which mutations induced white hair, microphthalmia and deafness in mice. The expression of *mi* gene was not affected by MSH stimulation. Using neural crest cells, Kubota Y from St. Marianna University in Kawasaki suggested that the stem cell factor might play a role in the development of c-kit positive cells from neural crest and that there was a critical time (Days 0-5) when the stem cell factor induce c-kit positive cells and that other factors such as cholera toxin might be necessary to induce further differentiation of the cells to melanocytes. Meyskens FL from University of California focused on the transcription factors of human melanocyte in response to ultraviolet radiation. His group demonstrated the high dose of 500 mJ of ultraviolet induced AP-1 and down regulated NF κ B and that normal human melanocytes required both the PKC and PKA signaling pathways for UVB induction of AP-1 and NF κ B. To elucidate the paracrine linkage of cytokines between human keratinocytes or fibroblasts and melanocytes for biological mechanisms of involved in cutaneous melanosis, Imokawa G from Kao Corporation characterized keratinocyte- and fibroblast-derived factors responsible for proliferation of melanocytes. They showed that IL-1 α and endothelin(ET)-1, and GM-SCF were

predominantly produced by keratinocytes in response to UVB and UVA irradiation, respectively, whereas stem cell factor (SCF) was the major cytokines produced by fibroblasts, and that these factors were synergistic of additive stimulatory effects on DNA synthesis of cultured melanocytes. Prota G from University of Naples suggested that melanin-related metabolites might play a critical role of the well being of both melanocytes and the surrounding cells. He showed that DHI was capable of inhibiting lipoxygenase-induced oxidation of arachidonic acid, a primary event in inflammatory reactions, and that DHI and DHICA were also endowed of excellent antioxidant properties and were capable of scavenging oxygen radicals. Another study was also presented that DHICA was a potent enhancer of nitric oxide production by LPS stimulated murine macrophages. To characterize melanins in mouse and human hair of various colors, Ito S from Fujita Health University School of Health Science presented new findings in their methods; 1) solubilization of hair melanins in Soluene-350 was a convenient method to estimate the total amount of melanin. 2) chemical differences among melanins produced in the brown, slaty, and other colored mice could be elucidated by HPLC and spectrophotometric methods. 3) eumelanin to pheomelanin ratio could be estimated from the slope of absorption spectrum of melanin dissolved in Soluene-350. Demonstrating the effects of prostanoids, leukotrienes, lymphokines, cytokines, melanotropins, endothelins and other factors on melanocyte on various conditions, Nordlund JJ from University of Cincinnati concluded that the concepts about the melanocyte and its receptors must be updated to show the multiplicity of receptors that made the cell responsible to many cytokines, lymphokines and chemical mediators of inflammation, and that the melanocyte itself had the same capabilities for forming these various factors as other cells, and that the formation of these factors by the melanocytes suggested that melanocytes had both autocrine and paracrine roles for all epidermal cells including Langerhans cells. Kikuchi K from University of Tokyo examined the ET receptor subtypes involved in mitogenic signaling in human primary and metastatic melanoma, and suggested that the mitogenic effects of ET in human primary melanoma were mainly dedicated through ETB receptors, and that down-regulation of ETB receptors caused the decreased growth response of ET-1 in metastatic melanoma cells.

2) Immunological Approaches for Malignant Melanoma

Itoh K from Kurume University showed that the MAGE-1 and -4 proteins which were tumor-rejection antigens recognized by cytotoxic T lymphocytes, were expressed in many different cancers including melanoma, and in spermatogonia and primary spermatocytes. And he suggested that MAGE proteins might be appropriate target molecules for specific immunotherapy of cancer. Ferrone S from New York Medical College Valhalla presented the data of immunotherapy with the anti-idiotypic (anti-id) antibodies. He discussed the rationale underlying the selection of anti-id mAb as immunogens and of human high molecular weight-melanoma associated antigen as a target, and described the immunogenic and structural characteristics of the anti-id mAb MK2-23. He also presented the results of active specific immunotherapy with mAb MK2-23 in 50 patients with advanced melanoma. To reduce the immunogenicity of anti-id mAb MK2-23, Matsumoto K from Shinshu University School of Medicine evaluated the *in vitro* reactivity and immunogenicity of F(ab')₂ fragments of mAb MK2-23 and of its chimeric form, and suggested that these might be useful immunogens to implement active specific immunotherapy in the patients with malignant melanoma. Hayashibe K from Kobe University School of Medicine analyzed a human melanoma-associated antigen D-1 which was identified by his group. They demonstrated that D-1 antigens were specifically expressed in melanoma cells and that HLA A33 might be associated with high expression group of D-1 antigen in patient with malignant melanoma. Taniguchi M from Chiba University investigated TCR repertoire in tumor-infiltrating lymphocytes in metastatic melanomas. He reported that only two TCR V α s, such as V α 3+ and V α 4+ TCRs, dominated and comprised about 75% of total TIL TCR V α repertoire, and that the majority of these two TCRs were extremely homogenous. And he also demonstrated that depletion of V α 3 T cells by anti-V α 3 resulted in protection against melanoma lung metastasis. Hearing VJ from National Institute of Health in Bethesda showed that many spontaneous autoimmune responses against melanocytes, including those directed against melanoma cells, reacted with epitopes derived from melanosomal proteins. He demonstrated that TRP2 was the most generally expressed of those potential antigens and thus might be the most appropriate target for vaccine strategies.

3) Biochemical Analysis on the Expression of Specific Proteins in Melanoma

Horikoshi T from Sapporo Medical University reported the serum 5-S-cysteinyl-dopa (5-S-CD) level reflects melanoma progression more sensitively than urinary 5-S-CD, serum or urinary DHICA, and

suggested that serum 5-S-CD might be the best biochemical marker for the detection of progression of melanotic melanoma. Melanin-producing cells are subject to a high risk of oxidative stress, particularly due to the presence of melanogenic machinery continuously producing o-quinones, the precursors of polymer melanin. Pavel S from Leiden University Hospital suggested that O-methylation was one of the protective means decreasing redox cycling potential of melanin precursors and speeding up their transmembrane transport, and that L-cysteine and glutathione not only participated in the redox reactions but also controlled the quality and quantity of produced melanin. Taniguchi S from Shinshu University of Medicine found and cloned a variant actin (β m actin) which was responsible for the decrease of metastatic ability of mouse melanoma cell. When the β m actin cDNA expression vector was transfected into a highly metastatic cell lines, he observed the inhibitory effects on the cell motility, invasion, and metastatic ability depending on the expression of the exogenously transferred β m actin. He suggested that β m actin inhibits the dynamic conversion between the monomeric and polymerized form of actin, leading to both a decrease in cell motility and consequently the suppression of invasiveness and metastasis. Kageshita T from Kumamoto University School of Medicine examined 58 primary and 35 metastatic melanoma and 22 pigmented nevi using anti-vitronectin receptor (VN-R) chain antibody, and found that VN-R avb3 chains were expressed in 47, 34 and one samples, respectively. He suggested that the expression of VN-R avb3 chain in melanocytic tumors was correlated with development of deep invasion and metastatic process. Tsuchida T from Saitama Medical School demonstrated that among gangliosides, GM3 and GD3 was predominant in congenital pigmented nevi and primary malignant melanoma, respectively. And he also reported that moderate amounts of sulfatide were detected in congenital pigmented nevi and primary malignant melanomas, but not in metastatic melanomas.

4) Diagnosis of Malignant Melanoma

Malignant melanoma has a variable prognosis determined by several specific histologic features. As the histopathologic variables, Mihm MC from Albany Medical College listed these parameters; 1. tumor thickness, 2. level invasion, 3. mitotic rate (per millimeter square), 4. tumor infiltrating lymphocytes, 5. regression, 6. ulceration, 7. predominant cell type morphology, 8. microscopic satellites, 9. vascular invasion, 10. nodular growth (vertical growth phase). Of these parameters, he emphasized tumor infiltrating lymphocytes and discussed the predominant cell morphology, its cell biologic implications as well as its molecular characteristics and how they related to prognosis. Hara H from Nihon University School presented several methods for the detection of melanogenic and proliferative activities on malignant blue nevus; formaldehyde-induced fluorescence from formalin-fixed, paraffin-embedded materials, HPLC analysis of 5-S-cysteinyldopa from frozen specimen, argyophilic nucleolar organizer regions (AgNORs) and cytofluorometry. He concluded that the cells on malignant blue nevus showed a melanogenic activity and that cytofluorometric analysis could be regarded as a useful parameter for the determination of proliferative activity. Investigating the videomicroscopic features of 500 melanocytic nevi on the soles, Akasu R from Yamanashi Medical University indicated that the surface profile of benign melanocytic nevi was classified into 5 types, and that malignant melanoma *in situ* and acral lentiginous melanoma were exclusively compartmentalized in the miscellaneous type. And she suggested that epiluminescence microscopy might be a useful method for discrimination of plantar benign and malignant melanocytic lesions and might be useful for long-term follow-up of the melanocytic lesions. Sober AJ from Harvard Medical School presented several new techniques in the early diagnosis of melanoma; an epiluminescence microscopy, a computerized image analysis and a confocal laser microscopy. Confocal laser microscopy utilizes an argon, krypton, or Ti:sapphire laser into a confocal scanning microscope producing high resolution images. His group was investigating the use of this technique in the imaging of pigmented lesions and assessing the clinicopathologic correlation. McCarthy WH from Royal Prince Alfred Hospital in Sydney used lymphoscintigraphy to mark the surface location of the sentinel lymph node in each node field and to measure the depth of the node, and demonstrated that lymphoscintigraphy was accurate in 97% of patients in identifying potential sites of micrometastases of sentinel nodes.

5) Treatment of Malignant Melanoma

Jimbow K from University of Alberta in Canada investigated the anti-melanoma effect of the phenolic thioether amines for the development of a new targeted chemotherapy of radiochemotherapy for malignant melanoma. He evaluated the specificity and improved effectiveness of N-propionyl cysteaminyphenol (NPrCAP) and its prodrug dipropionyl CAP (DPrCAP) over N-acetyl CAP (NAcCAP) and its prodrug of diacetyl form (DPrCAP) as antimelanoma agents, and the mechanism of

drug action of NAcCAP and NPrCAP focusing on the production of quinone/semiquinone radicals mediated through interaction with tyrosinase and other oxidases. Cascinelli N from National Cancer Institute in Milano discussed three specific and actual subjects of high interest: adoption of a new technique for an earlier detection of nodal metastases, gene therapy of metastatic disease and interferon in the adjuvant therapy of patients with nodal metastases. Riley PA from University College London Medical School tested sixteen phenolic compounds for their ability to act as substrates for tyrosinase and their cytotoxic potentials. He revealed that protection of the hydroxy group by acetylation or glycosylation prevented *in vitro* oxidation by tyrosinase, and that both acetate and succinate esters were cytotoxic, and that protection of the phenolic OH group by sugars abolished *in vitro* cytotoxicity. Isolated limb perfusion (ILP) with melphalan produces a 50% complete remission rate in melanoma and a 6% in sarcoma of the limbs. In order to improve these results, Lejeune FJ from Universitaire Vandois in Switzerland added rTNF α to melphalan and rIFN- γ in ILP for in-transit melanoma metastases, irresectable soft-tissue sarcomas and carcinomas of the limbs. A 90% complete response rate was obtained for melanoma, a 36.4% complete response rate for sarcoma and a 57% complete response rate for squamous cell carcinoma. Thioureylenes such as 2-thiouracil are known to be selectively accumulated in nascent melanin. On animal experiments, Larsson BS from Uppsala University in Sweden showed that the thioureylenes might be used as vehicles of radionuclides for melanoma scanning or treatment, and of boron-10 for neutron capture therapy. He also indicated that radioiodinated 2-thiouracil might be useful in the diagnosis of disseminated malignant melanoma. Hyperthermic isolated limb perfusion with the infusion of chemotherapeutic agents and/or cytokines has been proven to be definitely effective for the treatment of malignant melanoma at least locoregionally. Nakayama J from Kyushu University studied the mechanism of cytotoxic effects of hyperthermia on melanoma cells, and indicated that hyperthermia caused activation of immune system of the host, probably through the upregulation of ICAM-1 expression in melanoma cell and/or immune competent cell of the host.

Bibliography :

The Bibliography published in this issue covers the period February, 1996 through April, 1996. If you notice a paper that was not detected by this search that should be included, please send it to us and we will include it in the next issue. We have attempted to highlight any publications which include a member of the PASPCR with a star.

MELANINS, MELANOGENS & MELANOGENESIS

- Agodi A, Stefani S, Corsaro C, Campanile F, Gribaldo S, Sichel G: Study of a melanic pigment of *Proteus mirabilis*. Res Microbiol 147:167-174 (1996).
- Broekhuysse RM, Winkens HJ, Kuhlmann ED: Intraperitoneally injected melanin is highly uveitogenic. Exp Eye Res 62:199-200 (1996).
- Brooks R: Melanin as a visual signal amplifier in male guppies. Naturwissenschaften 83:39-41 (1996).
- Carney JA, Ferreiro JA: The epithelioid blue nevus: A multicentric familial tumor with important associations, including cardiac myxoma and psammomatous melanotic schwannoma. Am J Surg Pathol 20:259-272 (1996).
- Chakraborty AK, Platt JT, Kim KK, Kwon BS, Bennet DC, Pawelek JM: Polymerization of 5,6-dihydroxyindole-2-carboxylic acid to melanin by the pmel 17/silver locus protein. Eur J Biochem 236:180-188 (1996).
- Corsaro C, Scalia M, Blanco AR, Aiello I, Sichel G: Melanins in physiological conditions protect against lipoperoxidation. A study on albino and pigmented *Xenopus*. Pigm Cell Res 8:279-282 (1995).
- Deldotto P, Colzi A, Pardini C, Lucetti C, Dubini A, Grimaldi R, Bonuccelli U: Cabergoline improves motor disability without modifying L-DOPA plasma levels in fluctuating Parkinson's disease patients. J Neural Transmission-Suppl 259-265 (1995).
- Dirocco A, Molinari SP, Kollmeier B, Yahr MD: Parkinson's disease: Progression and mortality in the L-DOPA era. Parkinson'S Disease 69:3-11 (1996).
- Eldred, Ohbayashi, Jolly: Hermansky-Pudlak syndrome; A case report with analysis of auto-fluorescent ceroid-like pigments - Discussion. Gerontology 41:303(1995).
- Eller MS, Ostrom K, Gilchrest BA: DNA damage enhances melanogenesis. Proc Natl Acad Sci USA 93:1087-1092 (1996).
- Gebarski SS, Blaivas MA: Imaging of normal leptomeningeal melanin. Amer J Neuroradiol 17:55-60 (1996).

- Goto S, Yin D, James TJ, Zs Nagy I, Tatariunas: Studies on age pigments evolving into a new theory of biological aging - Discussion. *Gerontology* 41:171-172 (1995).
- Govind S: Rel signalling pathway and the melanotic tumour phenotype of *Drosophila*. *Biochem Soc Trans* 24:39-44 (1996).
- Hatano K, Ishiwata K, Yanagisawa T: Co production of 2,6-[F¹⁸] difluoroDOPA during electrophilic synthesis of 6-[F¹⁸]fluoro-L DOPA. *Nucl Med Biol* 23:101-103 (1996).
- Havaux M, Tardy F: Temperature-dependent adjustment of the thermal stability of photosystem II *in vivo*: Possible involvement of xanthophyll-cycle pigments. *Planta* 198:324-333 (1996).
- Hervieu G, Volant K, Grishina O, Descroixvagne M, Nahon JL: Similarities in cellular expression and functions of melanin-concentrating hormone and atrial natriuretic factor in the rat digestive tract. *Endocrinology* 137:561-571 (1996).
- Hoting E, Zimmermann M, Hocker H: Photochemical alterations in human hair .2. Analysis of melanin. *J Soc Cosmet Chem* 46:181-190 (1995).
- Ikeda K, Masujima T, Suzuki K, Sugiyama M: Cloning and sequence analysis of the highly expressed melanin-synthesizing gene operon from *Streptomyces castaneoglobosporus*. *Appl Microbiol Biotechnol* 45:80-85 (1996).
- Imada T, Kijima H, Takeuchi M, Shinkai S: Selective binding of glucose-6-phosphate, 3,4-dihydroxyphenylalanine (DOPA) and their analogs with a boronic-acid-appended metalloporphyrin. *Tetrahedron* 52:2817-2826 (1996).
- Imokawa G, Yada Y, Kimura M: Signalling mechanisms of endothelin-induced mitogenesis and melanogenesis in human melanocytes. *Biochem J* 314:305-312 (1996).
- Ishikawa T, Dhawan V, Chalhy T, Margouleff C, Robeson W, Dahl JR, Mandel F, Spetsieris P, Eidelberg D: Clinical significance of striatal DOPA decarboxylase activity in Parkinson's disease. *J Nucl Med* 37:216-222 (1996).
- Kingsmith C, Chen P, Garcia D, Rey H, Burnside B: Calcium-independent regulation of pigment granule aggregation and dispersion in teleost retinal pigment epithelial cells. *J Cell Sci* 109:33-43 (1996).
- Matsuda H, Higashino M, Nakai Y, Iinuma M, Kubo M, Lang FA: Studies of cuticle drugs from natural sources .4. Inhibitory effects of some arctostaphylos plants on melanin biosynthesis. *Biol Pharm Bull* 19:153-156 (1996).
- McClintock TS, Rising JP, Lerner MR: Melanophore pigment dispersion responses to agonists show two patterns of sensitivity to inhibitors of cAMP- dependent protein kinase and protein kinase C. *J Cell Physiol* 167:1-7 (1996).
- Meier H, Dullweber U: Bis(stilbenyl)squaraines - Novel pigments with extended conjugation. *Tetrahedron Lett* 37:1191-1194 (1996).
- Nappi AJ, Vass E, Prota G, Memoli S: The effects of hydroxyl radical attack on dopa, dopamine, 6-hydroxydopa, and 6-hydroxydopamine. *Pigm Cell Res* 8:283-293 (1995).
- Nutt JG: Effects of catechol-O-methyltransferase (COMT) inhibition on the pharmacokinetics of L-DOPA. *Parkinson'S Disease* 69:493-496 (1996).
- ☞ Oka M, Ichihashi M, Chakraborty AK: Enhanced expression of protein kinase C subspecies in melanogenic compartments B16 melanoma cells by UVB or MSH. *J Invest Dermatol* 106:377-378 (1996).
- Parkes DG: Diuretic and natriuretic actions of melanin concentrating hormone in conscious sheep. *J Neuroendocrinol* 8:57-63 (1996).
- ☞ Potterf SB, Muller J, Bernardini I, Tietze F, Kobayashi T, Hearing VJ, Gahl WA: Characterization of a melanosomal transport system in murine melanocytes mediating entry of the melanogenic substrate tyrosine. *J Biol Chem* 271:4002-4008 (1996).
- Prutz WA, Butler J, Land EJ: Photocatalytic and free radical interactions of the heterocyclic N-oxide resazurin with NADH, GSH and dopa. *Arch Biochem Biophys* 327:239-248 (1996).
- Reztsova NA, Kulish MA, Mironov AF: Plasma desorption mass spectrometry of natural substances .2. Applications of the method to the analysis of nucleotides, carbohydrates, lipids, and pigments. *Bioorg Khim* 22:83-93 (1996).
- Sagelsdorff P, Haenggi R, Heuberger B, Joppichkuhn R, Jung R, Weideli HJ, Joppich M: Lack of bioavailability of dichlorobenzidine from diarylide azo pigments: Molecular dosimetry for hemoglobin and DNA adducts. *Carcinogenesis* 17:507-514 (1996).
- Schmidt R, Krien P, Regnier M: The use of diethylaminoethyl-cellulose membrane filters in a bioassay to quantify melanin synthesis. *Anal Biochem* 235:113-118 (1996).
- Strid A, Chow WS, Anderson JM: Changes in the relaxation of electrochromic shifts of photosynthetic pigments and in the levels of mRNA transcripts in leaves of *Pisum sativum* as a result of exposure to supplementary UV-B radiation. The dependency on the intensity of the photosynthetically active radiation. *Plant Cell Physiol* 37:61-67 (1996).
- ☞ Sugumaran M: Oxidation of 3,4-dihydroxybenzylamine affords 3,4-dihydroxybenzaldehyde via the quinone methide intermediate. *Pigm Cell Res* 8:250-254 (1995).
- Suiko M, Sakakibara Y, Nakajima H, Sakaida H, Liu MC: Enzymic sulphation of dopa and tyrosine isomers by HepG2 human hepatoma cells: Stereoselectivity and stimulation by Mn²⁺. *Biochem J* 314:151-158 (1996).
- Thines E, Daussmann T, Semar M, Sterner O, Anke H: Fungal melanin biosynthesis inhibitors: Introduction of a test system based on the production of dihydroxynaphthalene (DHN) melanin in agar cultures. *Z Naturforsch C* 50:813-819 (1995).
- Tsai MJ, Lee EHY: Characterization of L-DOPA transport in cultured rat and mouse astrocytes. *J Neurosci Res* 43:490-495 (1996).
- Wang YL, Casadevall A: Susceptibility of melanized and nonmelanized *Cryptococcus neoformans* to the melanin-binding compounds trifluoperazine and chloroquine. *Antimicrob Agents Chemother* 40:541-545 (1996).

- Yin DZ: Studies on age pigments evolving into a new theory of biological aging. *Gerontology* 41:159-170 (1995).
- ♀ Zhao HQ, Eling DJ, Medrano EE, Boissy RE: Retroviral infection with human tyrosinase-related protein-1 (TRP-1) cDNA upregulates tyrosinase activity and melanin synthesis in a TRP-1-deficient melanoma cell line. *J Invest Dermatol* 106:744-752 (1996).
- Zoldan J, Friedberg G, Weizman A, Melamed E: Ondansetron, a 5-HT₃ antagonist for visual hallucinations and paranoid delusional disorder associated with chronic L-DOPA therapy in advanced Parkinson's disease. *Parkinson'S Disease* 69:541-544 (1996).

MELANOCYTES & KERATINOCYTES

- Abel PL, Olavarria JF: The callosal pattern in striate cortex is more patchy in monocularly enucleated albino than pigmented rats. *Neurosci Lett* 204:169-172 (1996).
- Allen MH, Robinson MK, Stephens PE, Macdonald DM, Barker JNWN: E-selectin binds to squamous cell carcinoma and keratinocyte cell lines. *J Invest Dermatol* 106:611-615 (1996).
- Beischer NA, Wein P: Linea alba pigmentation and umbilical deviation in nulliparous pregnancy: The ligamentum teres sign. *Obstet Gynecol* 87:254-256 (1996).
- Bessou S, Surlevebazeille JE, Sorbier E, Taieb A: *Ex vivo* reconstruction of the epidermis with melanocytes and the influence of UVB. *Pigm Cell Res* 8:241-249 (1995).
- Biolo R, Jori G, Soncin M, Rihter B, Kenney ME, Rodgers MAJ: Effect of photosensitizer delivery system and irradiation parameters on the efficiency of photodynamic therapy of B16 pigmented melanoma in mice. *Photochem Photobiol* 63:224-228 (1996).
- ♀ Boissy RE, Lamoreux ML: *In vivo* and *in vitro* morphological analysis of melanocytes homozygous for the mi(sp) allele at the murine *microphthalmia* locus. *Pigm Cell Res* 8:294-301 (1995).
- Bravo SM, Winalski CS, Weissman BN: Pigmented villonodular synovitis. *Radiol Clin North Am* 34:311(1996).
- Delmarmol V, Beermann F: Tyrosinase and related proteins in mammalian pigmentation. *FEBS Lett* 381:165-168 (1996).
- Deraeve LE, Deconinck AL, Dierichx PR, Roseeuw DI: Neonatal curettage of giant congenital melanocytic nevi. *Arch Dermatol* 132:20-22 (1996).
- Devries TJ, Verheijen JH, Debart ACW, Weidle UH, Ruiters DJ, Vanmuijen GNP: Decreased expression of both the low-density lipoprotein receptor-related protein $\alpha(2)$ -macroglobulin receptor and its receptor-associated protein in late stages of cutaneous melanocytic tumor progression. *Cancer Res* 56:1432-1439 (1996).
- Escobar C, Zuasti A, Ferrer C, Garciaortega F: Melanocytes in the stria vascularis and vestibular labyrinth of the mongolian gerbil (*Meriones unguiculatus*). *Pigm Cell Res* 8:271-278 (1995).
- ♀ Gahl WA, Potterf B, Durhampierre D, Brilliant MH, Hearing VJ: Melanosomal tyrosine transport in normal and pink-eyed dilution murine melanocytes. *Pigm Cell Res* 8:229-233 (1995).
- Giannini C, Scheithauer BW, Wenger DE, Unni KK: Pigmented villonodular synovitis of the spine: A clinical, radiological, and morphological study of 12 cases. *J Neurosurg* 84:592-597 (1996).
- Gilchrest BA, Park HY, Eller MS, Yaar M: Mechanisms of ultraviolet light-induced pigmentation. *Photochem Photobiol* 63:1-10 (1996).
- Grevelink JM, Duke D, Vanleeuwen RL, Gonzalez E, Decoste SD, Anderson RR: Laser treatment of tattoos in darkly pigmented patients: Efficacy and side effects. *J Am Acad Dermatol* 34:653-656 (1996).
- Hancock JA, Rivers JK, Mclean DI, Le ND, Gallagher RP: Melanocytic nevus density in Canadian native American Indian children. *J Am Acad Dermatol* 34:517-519 (1996).
- Herreid PA, Shapiro PE: Age distribution of spitz nevus vs malignant melanoma. *Arch Dermatol* 132:352-353 (1996).
- ♀ Horikawa T, Norris DA, Johnson TW, Zekman T, Dunscomb N, Bennion SD, Jackson RL, Morelli JG: DOPA-negative melanocytes in the outer root sheath of human hair follicles express premelanosomal antigens but not a melanosomal antigen or the melanosome-associated glycoproteins tyrosinase, TRP-1, and TRP-2. *J Invest Dermatol* 106:28-35 (1996).
- ♀ Huijzer JC, Mcfarland M, Niles RM, Meadows GG: Phorbol 12-myristate 13-acetate enhances *nm23* gene expression in murine melanocytes but not in syngeneic B16-BL6 melanoma variants. *J Cell Physiol* 166:487-494 (1996).
- Hunt G, Todd C, Thody AJ: Unresponsiveness of human epidermal melanocytes to melanocyte-stimulating hormone and its association with red hair. *Mol Cell Endocrinol* 116:131-136 (1996).
- Kay RM, Eckardt JJ, Mirra JM: Multifocal pigmented villonodular synovitis in a child: A case report. *Clin Orthop* 194-197 (1996).
- Kusch I, Markus M: Mollusc shell pigmentation: Cellular automaton simulations and evidence for undecidability. *J Theor Biol* 178:333-340 (1996).
- Maeda K, Fukuda M: Arbutin: Mechanism of its depigmenting action in human melanocyte culture. *J Pharmacol Exp Ther* 276:765-769 (1996).
- ♀ Marghoob AA, Schoenbach SP, Kopf AW, Orlov SJ, Nossa R, Bart RS: Large congenital melanocytic nevi and the risk for the development of malignant melanoma: A prospective study. *Arch Dermatol* 132:170-175 (1996).
- Meyer LJ, Piepkorn M, Goldgar DE, Lewis CM, Cannonalbright LA, Zone JJ, Skolnick MH: Interobserver concordance in discriminating clinical atypia of melanocytic nevi, and correlations with histologic atypia. *J Am Acad Dermatol* 34:618-625 (1996).

- Mitchell CH, Jacob TJC: A nonselective high conductance channel in bovine pigmented ciliary epithelial cells. *J Membrane Biol* 150:105-111 (1996).
- Monos T, Rosen SD, Karplus M, Yassur Y: Fundus pigmentation in retinopathy of prematurity. *Pediatrics* 97:343-348 (1996).
- Nakazawa K, Nakazawa H, Bonnard M, Damour O, Collombel C: Ca²⁺ and UVB radiation have no effect on E-cadherin-mediated melanocyte-keratinocyte adhesion. *Pigm Cell Res* 8:255-262 (1995).
- Naldi L, Adamoli L, Fraschini D, Corbetta A, Imberti L, Reseghetti A, Reciputo A, Rossi E, Cainelli T, Masera G: Number and distribution of melanocytic nevi in individuals with a history of childhood leukemia. *Cancer* 77:1402-1408 (1996).
- Norris A, Todd C, Graham A, Quinn AG, Thody AJ: The expression of the c-kit receptor by epidermal melanocytes may be reduced in vitiligo. *Br J Dermatol* 134:299-306 (1996).
- Numata K, Hiraki S, Yano H, Hayasaka S: Intracameral ascorbic acid, glutathione and protein levels in albino and pigmented rabbits. *Ophthalmic Res* 27:347-349 (1995).
- Ren ZP, Hedrum A, Ponten F, Nister M, Ahmadian A, Lundeberg J, Uhlen M, Ponten J: Human epidermal cancer and accompanying precursors have identical p53 mutations different from p53 mutations in adjacent areas of clonally expanded non-neoplastic keratinocytes. *Oncogene* 12:765-773 (1996).
- Schallreuter-Wood KU, Pittelkow MR, Swanson NN: Defective calcium transport in vitiliginous melanocytes. *Arch Dermatol Res* 288:11-13 (1996).
- Shanks JH, Banerjee SS: VS38 immunostaining in melanocytic lesions. *J Clin Pathol* 49:205-207 (1996).
- Shapiro S, Kessler S, Mcmenomey SO: Pathologic quiz case 1 - Pathologic diagnosis: Temporomandibular pigmented villonodular synovitis. *Arch Otolaryngol Head Neck Su* 122:194(1996).
- Stanganelli I, Rafanelli S, Bucchi L: Seasonal prevalence of digital epiluminescence microscopy patterns in acquired melanocytic nevi. *J Am Acad Dermatol* 34:460-464 (1996).
- Thody AJ: Epidermal melanocytes: Their regulation and role in skin pigmentation (vol 5, pg 558, 1995). *European J Dermatology* 6:155(1996).
- Trick CG, Wilhelm SW, Brown CM: Alterations in cell pigmentation, protein expression, and photosynthetic capacity of the cyanobacterium *Oscillatoria tenuis* grown under low iron conditions. *Can J Microbiol* 41:1117-1123 (1995).
- Walter MF, Zeineh LL, Black BC, Mcivor WE, Wright TRF, Biessmann H: Catecholamine metabolism and *in vitro* induction of premature cuticle melanization in wild type and pigmentation mutants of *Drosophila melanogaster*. *Arch Insect Biochem Physiol* 31:219-233 (1996).
- Wang YP, Becker D: Differential expression of the cyclin-dependent kinase inhibitors p16 and p21 in the human melanocytic system. *Oncogene* 12:1069-1075 (1996).
- Wintzen M, Yaar M, Burbach JPH, Gilchrist BA: Proopiomelanocortin gene product regulation in keratinocytes. *J Invest Dermatol* 106:673-678 (1996).
- Young LHY, Howard MA, Hu LK, Kim RY, Gragoudas ES: Photodynamic therapy of pigmented choroidal melanomas using a liposomal preparation of benzoporphyrin derivative. *Arch Ophthalmol* 114:186-192 (1996).

MELANOMA & METASTASIS

- Albert DM, Ryan LM, Borden EC: Metastatic ocular and cutaneous melanoma: A comparison of patient characteristics and prognosis. *Arch Ophthalmol* 114:107-108 (1996).
- Albertini JJ, Cruse CW, Rapaport D, Wells K, Ross M, Deconti R, Berman CG, Jared K, Messina J, Lyman G, Glass F, Fenske N, Reintgen DS: Intraoperative radiolymphoscintigraphy improves sentinel lymph node identification for patients with melanoma. *Ann Surg* 223:217-224 (1996).
- Aznavoorian S, Stracke ML, Parsons J, Mcclanahan J, Liotta LA: Integrin $\alpha(v)\beta(3)$, mediates chemotactic and haptotactic motility in human melanoma cells through different signaling pathways. *J Biol Chem* 271:3247-3254 (1996).
- Bae I, Smith ML, Sheikh MS, Zhan QM, Scudiero DA, Friend SH, Oconnor PM, Fornace AJ: An abnormality in the p53 pathway following γ -irradiation in many wild-type p53 human melanoma lines. *Cancer Res* 56:840-847 (1996).
- Balch CM, Buzaid AC: Finally, a successful adjuvant therapy for high-risk melanoma. *J Clin Oncol* 14:1-3 (1996).
- Banfield CC, Dawber RPR: Nail apparatus melanoma. *J Am Acad Dermatol* 34:322(1996).
- Bartoli C, Bono A, Clemente C, Delprato I, Zurrida S, Cascinelli N: Clinical diagnosis and therapy of cutaneous melanoma *in situ*. *Cancer* 77:888-892 (1996).
- Becker JC, Pancook JD, Gillies SD, Mendelsohn J, Reisfeld RA: Eradication of human hepatic and pulmonary melanoma metastases in SCID mice by antibody-interleukin 2 fusion proteins. *Proc Natl Acad Sci USA* 93:2702-2707 (1996).
- Bergman W, Gruis N: Familial melanoma and pancreatic cancer. *N Engl J Med* 334:471(1996).
- Berrutti LE, McClain SA: Neurotropic desmoplastic melanoma. *J Histotechnol* 19:69-77 (1996).
- Biesterfeld S, Kusche M, Viereck E, Fuzesi L: Limited value of the NK1/C3-antibody for the differential diagnosis of Paget's disease of the nipple and intra-epidermal malignant melanoma. *Histopathology* 28:269-270 (1996).
- Bird SJ, Brown MJ, Shy ME, Scherer SS: Chronic inflammatory demyelinating polyneuropathy associated with malignant melanoma. *Neurology* 46:822-824 (1996).

- Blangero J, Tissot RG, Beattie CW, Amoss MS: Genetic determinants of cutaneous malignant melanoma in Sinclair swine. *Br J Cancer* 73:667-671 (1996).
- Blend MJ, Hyun H, Patel B, Sullivan K, Salk D: Radioimmunoscinigraphy in patients with early stage cutaneous malignant melanoma. *J Nucl Med* 37:252-257 (1996).
- Blochdaum B, Muller M, Meisinger V, Eichler HG, Fassolt A, Pehamberger H: Measurement of extracellular fluid carboplatin kinetics in melanoma metastases with microdialysis. *Br J Cancer* 73:920-924 (1996).
- Boon T, Vanderbruggen P: Human tumor antigens recognized by T lymphocytes. *J Exp Med* 183:725-729 (1996).
- Bosserhoff AK, Hein R, Bogdahn U, Buettner R: Structure and promoter analysis of the gene encoding the human melanoma-inhibiting protein MIA. *J Biol Chem* 271:490-495 (1996).
- Bottasso B, Mari D, Coppola R, Santoro N, Vaglini M, Mannucci PM: Hypercoagulability and hyperfibrinolysis in patients with melanoma. *Thromb Res* 81:345-352 (1996).
- Bourke JF, Healsmith MF, Grahambrown RAC: Melanoma awareness and sun exposure in Leicester - Reply. *Br J Dermatol* 134:372-373 (1996).
- Brandt TP: Desmoplastic malignant melanoma. *J Am Acad Dermatol* 34:712-713 (1996).
- Brasoveanu LI, Alfomonte M, Fonsatti E, Colizzi F, Coral S, Nicotra MR, Cattarossi I, Cattelan A, Natali PG, Maio M: Levels of cell membrane CD59 regulate the extent of complement-mediated lysis of human melanoma cells. *Lab Invest* 74:33-42 (1996).
- Campaign JA, Slovak ML, Schoenlein PV, Popescu NC, Gottesman MM, Pastan I: Acquisition of multiple copies of a mutant topoisomerase II α allele by chromosome 17 aneuploidy is associated with etoposide resistance in human melanoma cell lines. *Somat Cell Mol Genet* 21:451-471 (1995).
- Celluzzi CM, Mayordomo JL, Storkus WJ, Lotze MT, Falo LD: Peptide-pulsed dendritic cells induce antigen-specific, CTL-mediated protective tumor immunity. *J Exp Med* 183:283-287 (1996).
- Char DH, Crawford JB, Kroll S: Iris melanomas: Diagnostic problems. *Ophthalmology* 103:251-255 (1996).
- Char DH, Kroll S, Quivey JM, Castro J: Long term visual outcome of radiated uveal melanomas in eyes eligible for randomisation to enucleation versus brachytherapy. *Br J Ophthalmol* 80:117-124 (1996).
- Chen RH, Gaos CM, Frazier OH: Complete resection of a right atrial intracavitary metastatic melanoma. *Ann Thorac Surg* 61:1255-1257 (1996).
- Chu YW, Seftor EA, Romer LH, Hendrix MJC: Experimental coexpression of vimentin and keratin intermediate filaments in human melanoma cells augments motility. *Am J Pathol* 148:63-69 (1996).
- Ciotti P, Strigini P, Bianchiscarra G: Familial melanoma and pancreatic cancer. *N Engl J Med* 334:469-470 (1996).
- Cottam DW, Corbitt RH, Gomez DE, Rees RC, Thorgeirsson UP: Alterations in endothelial cell proteinase and inhibitor polarized secretion following treatment with interleukin-1, phorbol ester, and human melanoma cell conditioned medium. *J Cell Biochem* 60:148-160 (1996).
- Cox NH, Aitchison TC, Sirel JM, Mackie RM: Comparison between lentigo maligna melanoma and other histogenetic types of malignant melanoma of the head and neck. *Br J Cancer* 73:940-944 (1996).
- Davis LS, Warren FM: Desmoplastic malignant melanoma - Reply. *J Am Acad Dermatol* 34:714(1996).
- Deboer DK, Schwartz HS, Thelman S, Reynolds VH: Heterogeneous survival rates for isolated skeletal metastases from melanoma. *Clin Orthop* 277-283 (1996).
- Degen WGJ, Weterman MAJ, Vangroningen JJM, Cornelissen IMAH, Lemmers JPWM, Agterbos MA, Vankessel AG, Swart GWM, Bloemers HPJ: Expression of *nma*, a novel gene, inversely correlates with the metastatic potential of human melanoma cell lines and xenografts. *Int J Cancer* 65:460-465 (1996).
- Dellian M, Richert C, Gamarra F, Goetz AE: Photodynamic eradication of amelanotic melanoma of the hamster with fast acting photosensitizers. *Int J Cancer* 65:246-248 (1996).
- Doi F, Chi DDJ, Charuworn BB, Conrad AJ, Russell J, Morton DL, Hoon DSB: Detection of β -human chorionic gonadotropin mRNA as a marker for cutaneous malignant melanoma. *Int J Cancer* 65:454-459 (1996).
- Eton O, Talpaz M, Lee KH, Rothberg JM, Brell JM, Benjamin RS: Phase II trial of recombinant human interleukin-2 and interferon- α -2a: Implications for the treatment of patients with metastatic melanoma. *Cancer* 77:893-899 (1996).
- Falkson CI: Treatment of metastatic malignant melanoma. *Anti-Cancer Drug* 6:709-716 (1995).
- Farrell DJ, Kashyap AP, Ashcroft T, Morrill GN: Primary malignant melanoma of the bronchus. *Thorax* 51:223-224 (1996).
- Fathallahshaykh HM, Zimmerman C, Morgan H, Rushing E, Schold SC, Unwin DH: Response of primary leptomeningeal melanoma to intrathecal recombinant interleukin-2: A case report. *Cancer* 77:1544-1550 (1996).
- Feldinghabermann B, Habermann R, Saldivar E, Ruggeri ZM: Role of β 3 integrins in melanoma cell adhesion to activated platelets under flow. *J Biol Chem* 271:5892-5900 (1996).
- Fricker J: Interleukin 7 on trial for melanoma treatment. *Mol Med Today* 2:2(1996).
- Fritschi L, Siemiatycki J: Melanoma and occupation: Results of a case-control study. *Occup Environ Medicine* 53:168-173 (1996).
- Fujimura Y, Ohno T, Siddique H, Lee L, Rao VN, Reddy ESP: The EWS-ATF-1 gene involved in malignant melanoma of soft parts with t(12;22) chromosome translocation, encodes a constitutive transcriptional activator. *Oncogene* 12:159-167 (1996).
- Gerard CM, Bruyns C, Delvaux A, Baudson N, Dargent JL, Goldman M, Velu T: Loss of tumorigenicity and increased immunogenicity induced by interleukin-10 gene transfer in B16 melanoma cells. *Hum Gene Ther* 7:23-31 (1996).
- Goetz DJ, Elsabban ME, Hammer DA, Pauli BU: Lu-ECAM-1-mediated adhesion of melanoma cells to endothelium under conditions of flow. *Int J Cancer* 65:192-199 (1996).

- Goldstein AM, Tucker MA: Familial melanoma and pancreatic cancer - Reply. *N Engl J Med* 334:471(1996).
- Gomez L, Rubio MP, Martin MT, Vazquez JJ, Idoate M, Pastoride G, Pestana A, Seizinger BR, Barnhill RL, Castresana JS: Chromosome 17 allelic loss and NF1-GRD mutations do not play a significant role as molecular mechanisms leading to melanoma tumorigenesis. *J Invest Dermatol* 106:432-436 (1996).
- Gradilone A, Greco C, Gazzaniga P, Vercillo R, Napolitano M, Gandini O, Mattei F, Frati L, Agliano AM: BAX gene expression in melanoma metastases. *J Invest Dermatol* 106:382(1996).
- Guilloux Y, Lucas S, Brichard VG, Vanpel A, Viret C, Deplaen E, Brasseur F, Lethe B, Jotereau F, Boon T: A peptide recognized by human cytolytic T lymphocytes on HLA-A2 melanomas is encoded by an intron sequence of the N-acetylglucosaminyltransferase V gene. *J Exp Med* 183:1173-1183 (1996).
- Haluska FG, Housman DE: Recent advances in the molecular genetics of malignant melanoma. *Cancer Surv* 25:277-292 (1995).
- Han DS, Pottincklemenceau C, Imro MA, Scudeletti M, Doucet C, Puppo F, Broutyboye D, Vedrenne J, Sahraoui Y, Brailly H, Poggi A, Jasmin C, Azzarone B, Indiveri F: IL2 triggers a tumor progression process in a melanoma cell line MELP, derived from a patient whose metastasis increased in size during IL2/INF α biotherapy. *Oncogene* 12:1015-1023 (1996).
- Hanuske AR, Catimel G, Aamdal S, Huinink WT, Paridaens R, Pavlidis N, Kaye SB, Tevelde A, Wanders J, Verweij J: Phase II clinical trials with rhizoxin in breast cancer and melanoma. *Br J Cancer* 73:397-399 (1996).
- Holder JE, Colloby PS, Fletcher A, Camp RDR: Amelanotic superficial spreading malignant melanoma mimicking Bowen's disease. *Br J Dermatol* 134:519-521 (1996).
- Hussussian CJ: Hot papers - Cancer genetics - Germline p16 mutations in familial melanoma, by C.J. Hussussian, J.P. Struewing, A.M. Goldstein, P.A.T. Higgins, D.S. Ally, M.D. Sheahan, W.H. Clark, M.A. Tucker, D.C. Dracopoli. *Scientist* 10:15(1996).
- Ishiguro T, Nakajima M, Naito M, Muto T, Tsuruo T: Identification of genes differentially expressed in B16 murine melanoma sublines with different metastatic potentials. *Cancer Res* 56:875-879 (1996).
- Isokangas OP, Muhonen T, Kajanti M, Pyrhonen S: Radiation therapy of intracranial malignant melanoma. *Radiother Oncol* 38:139-144 (1996).
- Itoh S, Hayashi H, Watanabe N, Kobayashi Y, Takii T, Onozaki K: Interleukin 1 (IL-1) production is not essential for acquired resistance of human A375 melanoma cells to anti-proliferative effect of IL-1. *Int J Cancer* 65:805-811 (1996).
- Jaeger E, Bernhard H, Romero P, Ringhoffer M, Arand M, Karbach J, Ilsemann C, Hagedorn M, Knuth A: Generation of cytotoxic T-cell responses with synthetic melanoma-associated peptides *in vivo*: Implications for tumor vaccines with melanoma-associated antigens. *Int J Cancer* 66:162-169 (1996).
- Janssen K, Kuntze J, Busse H, Schmid KW: p53 oncoprotein overexpression in choroidal melanoma. *Modern Pathol* 9:267-272 (1996).
- Jean D, Bareli M, Huang SY, Xie KP, Rodrigueslima F, Hermann J, Frade R: A cysteine proteinase, which cleaves human C3, the third component of complement, is involved in tumorigenicity and metastasis of human melanoma. *Cancer Res* 56:254-258 (1996).
- Johnson TM: Current therapy for cutaneous melanoma - Reply. *J Am Acad Dermatol* 34:323(1996).
- Johnson TM: Nail apparatus melanoma - Reply. *J Am Acad Dermatol* 34:322(1996).
- Karakousis CP, Driscoll DL: Positive deep nodes in the groin and survival in malignant melanoma. *Am J Surg* 171:421-422 (1996).
- Karlsson M, Boeryd B, Carstensen J, Franlund B, Gustafsson B, Kagedal B, Sun XF, Wingren S: Correlations of Ki-67 and PCNA to DNA ploidy, S-phase fraction and survival in uveal melanoma. *Eur J Cancer* 32A:357-362 (1996).
- Karlsson M, Jungnelius U, Aamdal S, Boeryd B, Carstensen J, Kagedal B, Westberg R, Wingren S: Correlation of DNA ploidy and S-phase fraction with chemotherapeutic response and survival in a randomized study of disseminated malignant melanoma. *Int J Cancer* 65:1-5 (1996).
- Kato T, Suetake T, Sugiyama Y, Tabata N, Tagami H: Epidemiology and prognosis of subungual melanoma in 34 Japanese patients. *Br J Dermatol* 134:383-387 (1996).
- Katz WE, Ferson PF, Lee RE, Killinger WA, Thompson ME, Gorcsan J: Metastatic malignant melanoma to the heart. *Circulation* 93:1066(1996).
- Khan SM, Collins RJ, Dunn IS: A melanoma cell line sensitive to expression of a fusion protein binding the retinoblastoma protein family pocket domain. *Exp Cell Res* 223:171-182 (1996).
- Kirkwood JM, Strawderman MH, Ernstoff MS, Smith TJ, Borden EC, Blum RH: Interferon α -2b adjuvant therapy of high-risk resected cutaneous melanoma: The Eastern Cooperative Oncology Group trial EST 1684. *J Clin Oncol* 14:7-17 (1996).
- Kirsch KH, Korradi Y, Johnson JP: Mader: A novel nuclear protein over expressed in human melanomas. *Oncogene* 12:963-971 (1996).
- Knutson JR, Iida J, Fields GB, Mccarthy JB: CD44/chondroitin sulfate proteoglycan and α 2 β 1 integrin mediate human melanoma cell migration on type IV collagen and invasion of basement membranes. *Mol Biol Cell* 7:383-396 (1996).
- Kojima S, Icho T, Mori H, Arai T: Enhancing potency of neopterin toward B-16 melanoma cell damage induced by UV-A irradiation and its possible application for skin tumor treatment. *Anticancer Res* 15:1975-1980 (1995).
- Krouse RS, Royal RE, Heywood G, Weintraub BD, White DE, Steinberg SM, Rosenberg SA, Schwartzentruber DJ: Thyroid dysfunction in 281 patients with metastatic melanoma or renal carcinoma treated with interleukin-2 alone. *J Immunother* 18:272-278 (1995).

- Lasek W, Wankowicz A, Kuc K, Feleszko W, Giermasz A, Jakobisiak M: Augmentation of antitumor efficacy by the combination of actinomycin D with tumor necrosis factor- α and interferon- γ on a melanoma model in mice. *Oncology* 53:31-37 (1996).
- Ledda MF, Adris S, Bover L, Bravo AL, Mordoh J, Podhajcer OL: The role of *SPARC* gene in the tumorigenic capacity of human melanoma cells. *Medicina-Buenos Aires* 56:51-54 (1996).
- Lee JÉ, Loflin PT, Laud PR, Lu M, Reveille JD, Lawlor DA: The human leukocyte antigen TAP2 gene defines the centromeric limit of melanoma susceptibility on chromosome 6p. *Tissue Antigens* 47:117-121 (1996).
- Leonetti C, Dagnano I, Lozupone F, Valentini A, Geiser T, Zon G, Calabretta B, Citro G, Zupi G: Antitumor effect of c-myc antisense phosphorothioate oligodeoxynucleotides on human melanoma cells *in vitro* and in mice. *J Nat Cancer Inst* 88:419-429 (1996).
- Lewis JJ, Houghton AN: Definition of tumor antigens suitable for vaccine construction. *Semin Cancer Biol* 6:321-327 (1995).
- Liao SK, Perng YP, Lee LA, Chang KSS, Lai GM, Wong E, Ho YS: Newly established MST-1 tumour cell line and tumour-infiltrating lymphocyte culture from a patient with soft tissue melanoma (clear cell sarcoma) and their potential applications to patient immunotherapy. *Eur J Cancer* 32A:346-356 (1996).
- Lupetti R, Mortarini R, Panceri P, Sensi M, Anichini A: Interaction with fibronectin regulates cytokine gene expression in human melanoma cells. *Int J Cancer* 66:110-116 (1996).
- Maloney JM: Current therapy for cutaneous melanoma. *J Am Acad Dermatol* 34:322-323 (1996).
- Marchand M, Weynants P, Rankin E, Arienti F, Belli F, Parmiani G, Cascinelli N, Bourlond A, Vanwijck R, Humblet Y, Canon JL, Laurent C, Naeyaert JM, Plagne R, Deraemaeker R, Knuth A, Jager E, Brasseur F, Herman J, Coulie PG, Boon T: Tumor regression responses in melanoma patients treated with a peptide encoded by gene *MAGE-3*. *Int J Cancer* 63:883-885 (1995).
- Marincola FM, Shamamian P, Rivoltini L, Salgaller M, Cormier J, Restifo NP, Simonis TB, Venzon D, White DE, Parkinson DR: HLA associations in the antitumor response against malignant melanoma. *J Immunother* 18:242-252 (1995).
- Martinezaguilan R, Seftor EA, Seftor REB, Chu YW, Gillies RJ, Hendrix MJC: Acidic pH enhances the invasive behavior of human melanoma cells. *Clin Exp Metastasis* 14:176-186 (1996).
- Mattavelli F, Dipalma S, Guzzo M: Primary mucosal malignant melanoma of the larynx: Case report and review of the literature. *Tumori* 81:460-463 (1995).
- Mccluggage WG, Shah V, Toner PG: Primary cutaneous myxoid malignant melanoma. *Histopathology* 28:179-182 (1996).
- Mcdermott VG, Low VHS, Keogan MT, Lawrence JA, Paulson EK: Malignant melanoma metastatic to the gastrointestinal tract. *Am J Roentgenol* 166:809-813 (1996).
- Mclean W, Gamel JW: Uveal (intraocular) melanoma. *Canc J* 9:20-25 (1996).
- Melia J, Chamberlain J: Melanoma awareness and sun exposure in Leicester. *Br J Dermatol* 134:372(1996).
- Menter DG, Fitzgerald L, Patton JT, Mcintire LV, Nicolson GL: Human melanoma integrins contribute to arrest and stabilization potential while flowing over extracellular matrix. *Immun Cell Biol* 73:575-583 (1995).
- Miele ME, Mccarty CT, Welch DR: SOD2 (MnSOD) does not suppress tumorigenicity or metastasis of human melanoma C8161 cells. *Anticancer Res* 15:2065-2070 (1995).
- Mihm MC, Clemente CG, Cascinelli N: Tumor infiltrating lymphocytes in lymph node melanoma metastases: A histopathologic prognostic indicator and an expression of local immune response. *Lab Invest* 74:43-47 (1996).
- Miller JM, Slater DN: Do histopathology reports of primary cutaneous melanoma contain enough essential information? *J Clin Pathol* 49:202-204 (1996).
- Miller N, Vile RG, Hart IR: Selectin ligands on human melanoma cells. *Glycoconjugate J* 13:33-43 (1996).
- Mossbacher U, Knollmayer S, Binder M, Steiner A, Wolff K, Pehamberger H: Increased nuclear volume in metastasizing "thick" melanomas. *J Invest Dermatol* 106:437-440 (1996).
- Muckenschnabel I, Bernhardt G, Spruss T, Buschauer A: Hyaluronidase pretreatment produces selective melphalan enrichment in malignant melanoma implanted in nude mice. *Cancer Chemother Pharmacol* 38:88-94 (1996).
- Mudun A, Murray DR, Herda SC, Eshima D, Shattuck LA, Vansant JP, Taylor AT, Alazraki NP: Early stage melanoma: Lymphoscintigraphy, reproducibility of sentinel node detection, and effectiveness of the intraoperative γ probe. *Radiology* 199:171-175 (1996).
- Mulder WMC, Stukart MJ, Roos M, Vanlier AW, Wagstaff J, Scheper RJ, Bloemena E: Culture of tumour-infiltrating lymphocytes from melanoma and colon carcinoma: Removal of tumour cells does not affect tumour-specificity. *Cancer Immunol Immunother* 41:293-301 (1995).
- Musk P, Clark JM, Thompson D, Dunn IS, Christopherson RI, Szabados E, Rose SE, Parsons PG: Purine deoxynucleoside metabolism in human melanoma cells with a high spontaneous mutation rate. *Mutat Res-Fundam Mol Mech Mut* 350:229-238 (1996).
- Nagatani T, Ichiyama S, Onuma R, Miyazawa M, Matsuzaki T, Miyagawa K, Baba N, Uchiyama M, Nakajima H: The use of DAV (DTIC, ACNU and VCR) and natural interferon- β combination therapy in malignant melanoma. *Acta Derm Venereol [Stockh]* 75:494(1995).
- Nazarian LN, Alexander AA, Rawool NM, Kurtz AB, Maguire HC, Mastrangelo MJ: Malignant melanoma: Impact of superficial US on management. *Radiology* 199:273-277 (1996).
- Niggli HJ: The cell nucleus of cultured melanoma cells as a source of ultraweak photon emission. *Naturwissenschaften* 83:41-44 (1996).

- Niles RM, Combs R: The relationship between susceptibility to retinoic acid treatment and protein kinase C α expression in murine melanoma cell lines. *Exp Cell Res* 223:20-28 (1996).
- Odonnell BF, Marsden JR, Odonnell CA, Sanders DSA, Billingham C: Does palpability of primary cutaneous melanoma predict dermal invasion? *J Am Acad Dermatol* 34:632-637 (1996).
- Ohta M, Berd D, Shimizu M, Nagai H, Cotticelli MG, Mastrangelo M, Shields JA, Shields CL, Croce CM, Huebner K: Deletion mapping of chromosome region 9p21-p22 surrounding the CDKN2 locus in melanoma. *Int J Cancer* 65:762-767 (1996).
- Ohta Y, Kijima H, Kashanisabet M, Scanlon KJ: Suppression of the malignant phenotype of melanoma cells by anti-oncogene ribozymes. *J Invest Dermatol* 106:275-280 (1996).
- Ohta Y, Kijima H, Ohkawa T, Kashanisabet M, Scanlon KJ: Tissue-specific expression of an anti-ras ribozyme inhibits proliferation of human malignant melanoma cells. *Nucleic Acids Res* 24:938-942 (1996).
- Orlandi L, Zaffaroni N, Bearzatto A, Costa A, Supino R, Vaglini M, Silvestrini R: Effect of melphalan and hyperthermia on cell cycle progression and cyclin B-1 expression in human melanoma cells. *Cell Proliferation* 28:617-630 (1995).
- Overgaard J, Gonzalez DG, Hulshof MCCH, Arcangeli G, Dahl O, Mella O, Bentzen SM: Hyperthermia as an adjuvant to radiation therapy of recurrent or metastatic malignant melanoma. A multicentre randomized trial by the European Society for Hyperthermic Oncology. *Int J Hyperthermia* 12:3-20 (1996).
- Papa MZ, Klein E, Karni T, Koller M, Davidson B, Azizi E, Benari G: Regional hyperthermic perfusion with cisplatin following surgery for malignant melanoma of the extremities. *Am J Surg* 171:416-420 (1996).
- Park SS, Walsh SM, Gragoudas ES: Visual-field deficits associated with proton beam irradiation for parapapillary choroidal melanoma. *Ophthalmology* 103:110-116 (1996).
- Penn I: Malignant melanoma in organ allograft recipients. *Transplantation* 61:274-278 (1996).
- Petit T, Borel C, Rixe O, Avril MF, Monnier A, Giroux B, Weil M, Khayat D: Complete remission seven years after treatment for metastatic malignant melanoma. *Cancer* 77:900-902 (1996).
- Pimm MV, Perkins AC, Strohal J, Ulbrich K, Duncan R: Gamma scintigraphy of the biodistribution of I¹²⁵-labelled N-(2-hydroxypropyl)methacrylamide copolymer-doxorubicin conjugates in mice with transplanted melanoma and mammary carcinoma. *J Drug Targeting* 3:375(1996).
- Platz A, Ringborg U, Lagerlof B, Lundqvist E, Sevigny P, Inganas M: Mutational analysis of the CDKN2 gene in metastases from patients with cutaneous malignant melanoma. *Br J Cancer* 73:344-348 (1996).
- Portner A, Etchison JR, Sampath D, Freeze HH: Human melanoma and Chinese hamster ovary cells galactosylate n-alkyl- β -glucosides using UDP gal:GlcNAc β 1,4galactosyltransferase. *Glycobiology* 6:7-13 (1996).
- Prehn RT: The paradoxical association of regression with a poor prognosis in melanoma contrasted with a good prognosis in keratoacanthoma. *Cancer Res* 56:937-940 (1996).
- Price EA, Coombe DR, Murray JC: Endothelial CD44H mediates adhesion of a melanoma cell line to quiescent human endothelial cells *in vitro*. *Int J Cancer* 65:513-518 (1996).
- Protti MP, Imro MA, Manfredi AA, Consogno G, Heltai S, Arcelloni C, Bellone M, Dellabona P, Casorati G, Rugarli C: Particulate naturally processed peptides prime a cytotoxic response against human melanoma *in vitro*. *Cancer Res* 56:1210-1213 (1996).
- Puk DE, Ketcham JM, Probst LE, Cameron JD, Holland EJ: Conjunctival malignant melanoma. *Arch Ophthalmol* 114:100-101 (1996).
- Reinhold U, Bruske T, Kreysel HW: Dacarbazine, vincristine, bleomycin and lomustine plus natural interferon- α for metastatic melanoma. *Eur J Cancer* 32A:180(1996).
- Restifo NP, Marincola FM, Kawakami Y, Taubenberger J, Yannelli JR, Rosenberg SA: Loss of functional β (2)-microglobulin in metastatic melanomas from five patients receiving immunotherapy. *J Nat Cancer Inst* 88:100-108 (1996).
- Riggs JE: Biphase Gompertzian malignant melanoma mortality rate dynamics: A manifestation of population genetic heterogeneity. *Ann Hum Biol* 22:515-523 (1995).
- Rivers JK: Melanoma. *Lancet* 347:803-806 (1996).
- Robbins PF, Elgamil M, Li YF, Kawakami Y, Loftus D, Appella E, Rosenberg SA: A mutated β -catenin gene encodes a melanoma-specific antigen recognized by tumor infiltrating lymphocytes. *J Exp Med* 183:1185-1192 (1996).
- Robinson ES, Dooley TP: A new allogeneic model for metastatic melanoma. *Eur J Cancer* 31A:2302-2308 (1995).
- Safarians S, Sternlicht MD, Freiman CJ, Huaman JA, Barsky SH: The primary tumor is the primary source of metastasis in a human melanoma scid model. Implications for the direct autocrine and paracrine epigenetic regulation of the metastatic process. *Int J Cancer* 66:151-158 (1996).
- Sagebiel RW: Desmoplastic malignant melanoma - Reply. *J Am Acad Dermatol* 34:713(1996).
- Sakamoto T, Sakamoto M, Yoshikawa H, Hata Y, Ishibashi T, Ohnishi Y, Inomata H: Histologic findings and prognosis of uveal malignant melanoma in Japanese patients. *Am J Ophthalmol* 121:276-283 (1996).
- Sander B, Boeryd B: Tumor necrosis factor- α expression in human primary malignant melanoma and its relationship to tumor infiltration by CD3(+) cells. *Int J Cancer* 66:42-47 (1996).
- Schwurzervoit M, Proebstle TM, Sterry W: Identification of lymph node metastases by use of polymerase chain reaction (PCR) in melanoma patients. *Eur J Cancer* 32A:264-268 (1996).
- Shields JA: Local resection of posterior uveal melanoma. *Br J Ophthalmol* 80:97-98 (1996).
- Singh AD, Shields CL, Shields JA, Depotter P: Bilateral primary uveal melanoma: Bad luck or bad genes? *Ophthalmology* 103:256-262 (1996).

- Sinha T, Benedict R: Relationship between latitude and melanoma incidence: International evidence. *Cancer Lett* 99:225-231 (1996).
- ☞ Soballe PW, Montone KT, Satyamoorthy K, Nesbit M, Herlyn M: Carcinogenesis in human skin grafted to SCID mice. *Cancer Res* 56:757-764 (1996).
- Stone CH, Lynch EF, Linden MD, Kubus JJ, Zarbo RJ: Immunocytochemical evaluation of proliferating cell nuclear antigen, Ki-67 (MIB-1), and p53 in predicting survival of primary and metastatic malignant melanomas. *Appl Immunohistochem* 4:25-33 (1996).
- Timar J, Liu B, Bazaz R, Honn KV: Association of protein kinase-C- α with cytoplasmic vesicles in melanoma cells. *J Histochem Cytochem* 44:177-182 (1996).
- Uhlenkott CE, Huijzer JC, Cardeiro DJ, Elstad CA, Meadows GG: Attachment, invasion, chemotaxis, and proteinase expression of B16-BL6 melanoma cells exhibiting a low metastatic phenotype after exposure to dietary restriction of tyrosine and phenylalanine. *Clin Exp Metastasis* 14:125-137 (1996).
- Urosevic V, Chollet P, Adenis A, Chauvergne J, Fargeot P, Roche H, Kerbrat P, Lentz MA, Fumoleau P, Chevallier B: Results of a phase II trial with cystemustine in advanced malignant melanoma. A trial of the EORTC clinical ScreeningGroup. *Eur J Cancer* 32A:181-182 (1996).
- Vangolen KL, Risin S, Staroselsky A, Berger D, Tainsky MA, Pathak S, Price JE: Predominance of the metastatic phenotype in hybrids formed by fusion of mouse and human melanoma clones. *Clin Exp Metastasis* 14:95-106 (1996).
- Viac J, Misery L, Schmitt D, Claudy A: Follow-up of circulating ICAM-1 in malignant melanoma: Correlation with the clinical course of the disease. *Br J Dermatol* 134:604-605 (1996).
- Villa R, Zaffaroni N, Bearzatto A, Costa A, Sichirillo A, Silvestrini R: Effect of ionizing radiation on cell-cycle progression and cyclin β 1 expression in human melanoma cells. *Int J Cancer* 66:104-109 (1996).
- Vilmer C, Bailly C, Ledoussal V, Lasry S, Guerin P, Delaunay MM, Mandard AM: Thin melanomas with unusual aggressive behavior: A report of nine cases. *J Am Acad Dermatol* 34:439-444 (1996).
- Wang M, Chen PW, Bronte V, Rosenberg SA, Restifo NP: Anti-tumor activity of cytotoxic T lymphocytes elicited with recombinant and synthetic forms of a model tumor-associated antigen. *J Immunother* 18:139-146 (1995).
- Wang RF, Parkhurst MR, Kawakami Y, Robbins PF, Rosenberg SA: Utilization of an alternative open reading frame of a normal gene in generating a novel human cancer antigen. *J Exp Med* 183:1131-1140 (1996).
- Wang YL, Uhara H, Yamazaki Y, Nikaido T, Saida T: Immunohistochemical detection of CDK4 and p16(INK4) proteins in cutaneous malignant melanoma. *Br J Dermatol* 134:269-275 (1996).
- Wassberg C, Thorn M, Yuen J, Ringborg U, Hakulinen T: Second primary cancers in patients with cutaneous malignant melanoma: A population-based study in Sweden. *Br J Cancer* 73:255-259 (1996).
- Whelan AJ, Bartsch D, Goodfellow PJ: Familial melanoma and pancreatic cancer - Reply. *N Engl J Med* 334:472(1996).
- White VA, Mcneil BK, Thiberville L, Horsman DE: Acquired homozygosity (isodisomy) of chromosome 3 during clonal evolution of a uveal melanoma: Association with morphologic heterogeneity. *Gene Chromosome Cancer* 15:138-143 (1996).
- Whiteman D, Valery P, Mcwhirter W, Green A: Incidence of cutaneous childhood melanoma in Queensland, Australia. *Int J Cancer* 63:765-768 (1995).
- Wright FA, Thomas RG: Familial melanoma and pancreatic cancer. *N Engl J Med* 334:470-471 (1996).
- ☞ Xu X, Azumi N, Skelton HG, Hearing VJ, Gersten DM: Immunostaining of human melanomas by a monoclonal antibody to B700 mouse melanoma antigen. *Eur J Cancer* 32A:168-173 (1996).
- Yamamura S, Sadahira Y, Ruan FQ, Hakomori S, Igarashi Y: Sphingosine-1-phosphate inhibits actin nucleation and pseudopodium formation to control cell motility of mouse melanoma cells. *FEBS Lett* 382:193-197 (1996).
- Yannelli JR, Mcconnell S, Parker L, Nishimura M, Robbins P, Yang J, Elgamil M, Kawakami Y: Melanoma tumor-infiltrating lymphocytes derived from four distinct anatomic sites obtained from a single patient: Comparison of functional reactivity and melanoma antigen recognition. *J Immunother* 18:263-271 (1995).
- Zehetmayer M, Menapace R, Kitz K, Ertl A: Gamma knife radiosurgery for uveal melanoma. *Radiosurgery* 1:165-171 (1995).
- Zong ZP, Fujikawayamamoto K, Tanino M, Teraoka K, Yamagishi H, Odashima S: Saikosaponin b(2)-induced apoptosis of cultured B16 melanoma cell line through down-regulation of PKC activity. *Biochem Biophys Res Commun* 219:480-485 (1996).

MSH, POMC, GROWTH FACTORS & RECEPTORS

- Adan RAH, Vanderkraan M, Doornbos RP, Bar PR, Burbach JPH: Melanocortin receptors mediate α -MSH-induced stimulation of neurite outgrowth in Neuro 2A cells. *Mol Brain Res* 36:37-44 (1996).
- Aguanno A, Afar R, Albert VR: Tissue-specific expression of the nonneuronal promoter of the aromatic L-amino acid decarboxylase gene is regulated by hepatocyte nuclear factor 1. *J Biol Chem* 271:4528-4538 (1996).
- Apte RS, Niederkorn JY: Isolation and characterization of a unique natural killer cell inhibitory factor present in the anterior chamber of the eye. *J Immunol* 156:2667-2673 (1996).
- Azzarone B, Pottinclenceau C, Krief P, Rubinstein E, Jasmin C, Scudeletti M, Indiveri F: Are interleukin-2 and interleukin-15 tumor promoting factors for human non-hematopoietic cells? *Eur Cytokine Netw* 7:27-36 (1996).

- Bellastella A, Sinisi AA, Criscuolo T, Debellis A, Carella C, Iorio S, Sinisi AM, Parlato F, Venditto T, Pisano G: Melatonin and the pituitary-thyroid axis status in blind adults: A possible resetting after puberty. *Clin Endocrinol* 43:707-711 (1995).
- Bernardi M, Bertolini A, Szczawinska K, Genedani S: Blockade of the polyamine site of NMDA receptors produces antinociception and enhances the effect of morphine, in mice. *Eur J Pharmacol* 298:51-55 (1996).
- Bhardwaj RS, Schwarz A, Becher E, Mahnke K, Aragane Y, Schwarz T, Luger TA: Pro-opiomelanocortin-derived peptides induce IL-10 production in human monocytes. *J Immunol* 156:2517-2521 (1996).
- Caputo FA, Ali SF, Wolff GL, Scallet AC: Neonatal MSG reduces hypothalamic DA, β -endorphin, and delays weight gain in genetically obese (A(viable yellow)/ a) mice. *Pharmacol Biochem Behav* 53:425-432 (1996).
- Chhajlani V: Characterization of a putative α -MSH antagonist 153N-6 at melanocortin receptor subtypes by radioligand binding. *Peptides* 17:349-351 (1996).
- Chhajlani V, Xu XL, Blauw J, Sudarshi S: Identification of ligand binding residues in extracellular loops of the melanocortin 1 receptor. *Biochem Biophys Res Commun* 219:521-525 (1996).
- Clark AJL, Cammas FM: The ACTH receptor. *Bailliere Clin Endocrinol Met* 10:29-47 (1996).
- Clemente CG, Mihm MG, Bufalino R, Zurrida S, Collini P, Cascinelli N: Prognostic value of tumor infiltrating lymphocytes in the vertical growth phase of primary cutaneous melanoma. *Cancer* 77:1303-1310 (1996).
- Collins VR, Dowse GK, Cabealawa S, Ram P, Zimmet PZ: High mortality from cardiovascular disease and analysis of risk factors in Indian and Melanesian Fijians. *Int J Epidemiol* 25:59-69 (1996).
- Damato BE, Paul J, Foulds WS: Risk factors for metastatic uveal melanoma after trans-scleral local resection. *Br J Ophthalmol* 80:109-116 (1996).
- Damato BE, Paul J, Foulds WS: Risk factors for residual and recurrent uveal melanoma after trans-scleral local resection. *Br J Ophthalmol* 80:102-108 (1996).
- Duckers HJ, Vandokkum RP, Verhaagen J, Dasilva FHL, Gispen WH: Functional and neurophysiological evidence of the efficacy of trophic pharmacotherapy using an adrenocorticotrophic hormone analog in experimental allergic encephalomyelitis, an animal model of multiple sclerosis. *Neuroscience* 71:507-521 (1996).
- Eaton MJ, Lookingland KJ, Moore KE: The sigma receptor ligand rimcazole alters secretion of prolactin and α -melanocyte stimulating hormone by dopaminergic and non-dopaminergic mechanisms. *Eur J Pharmacol* 299:171-177 (1996).
- Evans SRT, Houghton AM, Schumaker L, Brenner RV, Buras RR, Davoodi F, Nauta RJ, Shabahang M: Vitamin D receptor and growth inhibition by 1,25-dihydroxyvitamin D-3 in human malignant melanoma cell lines. *J Surg Res* 61:127-133 (1996).
- Fraker DL, Alexander HR, Andrich M, Rosenberg SA: Treatment of patients with melanoma of the extremity using hyperthermic isolated limb perfusion with melphalan, tumor necrosis factor, and interferon γ : Results of a tumor necrosis factor dose-escalation study. *J Clin Oncol* 14:479-489 (1996).
- Fritz KL, Koziol S, Fabian DF, Lefor AT: Tumor necrosis factor α mediates the antitumor effect of combined interleukin-2 and whole body hyperthermia. *J Surg Res* 60:55-60 (1996).
- Fujisawa T, Ikegami H, Yamato E, Takekawa K, Nakagawa Y, Hamada Y, Oga T, Ueda H, Shintani M, Fukuda M, Ogihara T: Association of Trp64Arg mutation of the β 3-adrenergic-receptor with NIDDM and body weight gain. *Diabetologia* 39:349-352 (1996).
- Fuse N, Yasumoto K, Suzuki H, Takahashi K, Shibahara S: Identification of a melanocyte-type promoter of the microphthalmia-associated transcription factor gene. *Biochem Biophys Res Commun* 219:702-707 (1996).
- Gage PJ, Roller ML, Saunders TL, Scarlett LM, Camper SA: Anterior pituitary cells defective in the cell-autonomous factor, *df*, undergo cell lineage specification but not expansion. *Development* 122:151-160 (1996).
- Garg PK, Alston KL, Welsh PC, Zalutsky MR: Enhanced binding and inertness to dehalogenation of α -melanotropic peptides labeled using N-succinimidyl 3-iodobenzoate. *Bioconjugate Chemistry* 7:233-239 (1996).
- Gelmann EP: Tamoxifen induction of apoptosis in estrogen receptor- negative cancers: New tricks for an old dog? *J Nat Cancer Inst* 88:224-226 (1996).
- Gonzalez M, Lezcano N, Celis ME, Fidelio GD: Interaction of α -MSH and substance P with interfaces containing gangliosides. *Peptides* 17:269-274 (1996).
- Gonzalez MI, Vaziri S, Wilson CA: Behavioral effects of α -MSH and MCH after central administration in the female rat. *Peptides* 17:171-177 (1996).
- Grabbe S, Bhardwaj RS, Mahnke K, Simon MM, Schwarz T, Luger TA: α -Melanocyte-stimulating hormone induces hapten-specific tolerance in mice. *J Immunol* 156:473-478 (1996).
- Griffioen AW, Damen CA, Martinotti S, Blijham GH, Groenewegen G: Endothelial intercellular adhesion molecule-1 expression is suppressed in human malignancies: The role of angiogenic factors. *Cancer Res* 56:1111-1117 (1996).
- Hamanoue M, Takemoto N, Matsumoto K, Nakamura T, Nakajima K, Kohsaka S: Neurotrophic effect of hepatocyte growth factor on central nervous system neurons *in vitro*. *J Neurosci Res* 43:554-564 (1996).
- Heaney ML, Golde DW: Soluble cytokine receptors. *Blood* 87:847-857 (1996).
- Hwang BJ, Liao JC, Chu G: Isolation of a cDNA encoding a UV-damaged DNA binding factor defective in xeroderma pigmentosum group E cells. *Mutat Res-DNA Repair* 362:105-117 (1996).
- Iveson RD, Guthrie DM, Veres DA, Doughty D: A study of the effect of anatomical site on sun protection factor efficiency using a novel UV delivery device. *J Soc Cosmet Chem* 46:271-280 (1995).

- Jiang HP, Lin JJ, Su ZZ, Goldstein NI, Fisher PB: Subtraction hybridization identifies a novel melanoma differentiation associated gene, mda-7, modulated during human melanoma differentiation, growth and progression. *Oncogene* 11:2477-2486 (1995).
- Kesavalu L, Holt SC, Ebersole JL: Trypsin-like protease activity of *Porphyromonas gingivalis* as a potential virulence factor in a murine lesion model. *Microb Pathog* 20:1-10 (1996).
- Kikuchi K, Nakagawa H, Kadono T, Etoh T, Byers HR, Mihm MC, Tamaki K: Decreased ET(B) receptor expression in human metastatic melanoma cells. *Biochem Biophys Res Commun* 219:734-739 (1996).
- Krystal GW, Hines SJ, Organ CP: Autocrine growth of small cell lung cancer mediated by coexpression of c-kit and stem cell factor. *Cancer Res* 56:370-376 (1996).
- Kumar V: Effects of melatonin in blocking the response to a skeleton photoperiod in the blackheaded bunting. *Physiol Behav* 59:617-620 (1996).
- Langtimmsedlak CJ, Schroeder B, Saskowski JL, Carnahan JF, Sieberblum M: Multiple actions of stem cell factor in neural crest cell differentiation *in vitro*. *Dev Biol* 174:345-359 (1996).
- Levraud JP, Pannetier C, Langladedemoyen P, Brichard V, Kourilsky P: Recurrent T cell receptor rearrangements in the cytotoxic T lymphocyte response *in vivo* against the P815 murine tumor. *J Exp Med* 183:439-449 (1996).
- Lezcano NE, Gonzalez M, Fidelio GD, Celis ME: Synergic action of gangliosides on α -MSH-induced cyclic AMP levels in rat brain slices. *Peptides* 17:345-347 (1996).
- Lichtensteiger W, Hanimann B, Siegrist W, Eberle AN: Region- and stage-specific patterns of melanocortin receptor ontogeny in rat central nervous system, cranial nerve ganglia and sympathetic ganglia. *Brain Res Dev Brain Res* 91:93-110 (1996).
- Longati P, Alberio D, Comoglio PM: Hepatocyte growth factor is a pleiotropic factor protecting epithelial cells from apoptosis. *Cell Death Differentiation* 3:23-28 (1996).
- Lugo DI, Pintar JE: Ontogeny of basal and regulated proopiomelanocortin- derived peptide secretion from fetal and neonatal pituitary intermediate lobe cells: Melanotrophs exhibit transient glucocorticoid responses during development. *Dev Biol* 173:110-118 (1996).
- Macedo ZS, Furquim TA, Ito AS: Estimation of average depth of penetration of melanotropins in dimyristoylphosphatidylglycerol vesicles. *Biophys Chem* 59:193-202 (1996).
- Maes M, Vangastel A, Meltzer HY, Cosyns P, Blockx P, Desnyder R: Acute administration of buspirone increases the escape of hypothalamic-pituitary-adrenal-axis hormones from suppression by dexamethasone in depression. *Psychoneuroendocrinology* 21:67-81 (1996).
- Martensson LGE, Andersson RGG: A melatonin binding site modulates the $\alpha(2)$ -adrenoceptor. *Life Sci* 58:525-533 (1996).
- Mattern J, Koomagi R, Volm M: Association of vascular endothelial growth factor expression with intratumoral microvessel density and tumour cell proliferation in human epidermoid lung carcinoma. *Br J Cancer* 73:931-934 (1996).
- Naville D, Barjhoux L, Jaillard C, Faury D, Despert F, Esteva B, Durand P, Saez JM, Begeot M: Demonstration by transfection studies that mutations in the adrenocorticotropin receptor gene are one cause of the hereditary syndrome of glucocorticoid deficiency. *J Clin Endocrinol Metab* 81:1442-1448 (1996).
- Nowicki A, Szenajch J, Ostrowska G, Wojtowicz A, Wojtowicz K, Kruszewski AA, Maruszynski M, Aukerman SL, Wiktorjedrzejczak W: Impaired tumor growth in colony-stimulating factor 1 (CSF- 1)-deficient, macrophage-deficient op/op mouse: Evidence for a role of CSF-1-dependent macrophages in formation of tumor stroma. *Int J Cancer* 65:112-119 (1996).
- Ohyama C, Kawamura S, Suzuki K, Numahata K, Tokuyama S, Ito A, Satoh M, Saito S, Yoshikawa K, Hoshi S, Orikasa S: G(M3) inhibits murine MET-2 tumor invasion and growth. *Int J Oncol* 8:809-813 (1996).
- Papadimitriou CA, Topp MS, Serve H, Oelmann E, Koenigsmann M, Maurer J, Oberberg D, Reufi B, Thiel E, Berdel WE: Recombinant human stem cell factor does exert minor stimulation of growth in small cell lung cancer and melanoma cell lines. *Eur J Cancer* 31A:2371-2378 (1995).
- Perr H, Oh P, Johnson D: Developmental regulation of transforming growth factor β -mediated collagen synthesis in human intestinal muscle cells. *Gastroenterology* 110:92-101 (1996).
- Polgar WE, Bitler CM, Toll L: Desensitization of melanotropin receptors in COS-7 cells. *Life Sci* 58:1223-1229 (1996).
- Presse F, Sorokovsky I, Max JP, Nicolaidis S, Nahon JL: Melanin-concentrating hormone is a potent anorectic peptide regulated by food-deprivation and glucopenia in the rat. *Neuroscience* 71:735-745 (1996).
- Qu DQ, Ludwig DS, Gammeltoft S, Piper M, Pelleymounter MA, Cullen MJ, Mathes WF, Przypek J, Kanarek R, Maratosflier E: A role for melanin-concentrating hormone in the central regulation of feeding behaviour. *Nature* 380:243-247 (1996).
- Rajora N, Ceriani G, Catania A, Star RA, Murphy MT, Lipton JM: α -MSH production, receptors, and influence on neopterin in a human monocyte macrophage cell line. *J Leukocyte Biol* 59:248-253 (1996).
- Rieber MS, Welch DR, Miele ME, Rieber M: p53-independent increase in p21WAF1 and reciprocal down-regulation of cyclin A and proliferating cell nuclear antigen in bromodeoxyuridine-mediated growth arrest of human melanoma cells. *Cell Growth Differ* 7:197-202 (1996).
- Robert C, Pouvelle B, Meyer P, Muanza K, Fujioka H, Aikawa M, Scherf A, Gysin J: Chondroitin-4-sulphate (proteoglycan), a receptor for *Plasmodium falciparum*-infected erythrocyte adherence on brain microvascular endothelial cells. *Res Immunol* 146:383-393 (1995).
- Robertson GP, Coleman AB, Lugo TG: Mechanisms of human melanoma cell growth and tumor suppression by chromosome 6(1). *Cancer Res* 56:1635-1641 (1996).

- Rosengren E, Bucala R, Aman P, Jacobsson L, Odh G, Metz CN, Rorsman H: The immunoregulatory mediator macrophage migration inhibitory factor (MIF) catalyzes a tautomerization reaction. *Mol Med* 2:143-149 (1996).
- Rothemberger S, Food MR, Gabathuler R, Kennard ML, Yamada T, Yasuhara O, Mcgeer PL, Jefferies WA: Coincident expression and distribution of melanotransferrin and transferrin receptor in human brain capillary endothelium. *Brain Res* 712:117-121 (1996).
- Siegrist W, Willard DH, Wilkison WO, Eberle AN: Agouti protein inhibits growth of B16 melanoma cells *in vitro* by acting through melanocortin receptors. *Biochem Biophys Res Commun* 218:171-175 (1996).
- Slominski A, Ermak G, Hwang J, Chakraborty A, Mazurkiewicz JE, Mihm M: Proopiomelanocortin, corticotropin releasing hormone and corticotropin releasing hormone receptor genes are expressed in human skin. *FEBS Lett* 374:113-116 (1996).
- Slominski A, Ermak G, Hwang J, Mazurkiewicz J, Corliss D, Eastman A: The expression of proopiomelanocortin (POMC) and of corticotropin releasing hormone receptor (CRH-R) genes in mouse skin. *Biochim Biophys Acta* 1289:247-251 (1996).
- Takahashi N, Fiers W, Brouckaert P: Anti-tumor activity of tumor necrosis factor in combination with interferon- γ is not affected by prior tolerization. *Int J Cancer* 63:846-854 (1995).
- Tosi E, Valota O, Canevari S, Adobati E, Casalini P, Perez P, Colnaghi MI: Anti-idiotypic response to antigrowth factor receptor monoclonal antibodies. *Eur J Cancer* 32A:498-505 (1996).
- Vanbergen P, Janssen PML, Hoogerhout P, Dewiltdt DJ, Versteeg DHG: Cardiovascular effects of γ -MSH ACTH-like peptides: Structure-activity relationship. *Eur J Pharmacol* 294:795-803 (1995).
- Vanstrien FJC, Jespersen S, Vandergreef J, Jenks BG, Roubos EW: Identification of POMC processing products in single melanotrope cells by matrix-assisted laser desorption/ ionization mass spectrometry. *FEBS Lett* 379:165-170 (1996).
- Viac J, Vincent C, Palacio S, Schmitt D, Claudy A: Tumour necrosis factor (TNF) soluble receptors in malignant melanoma: Correlation with soluble ICAM-1 levels. *Eur J Cancer* 32A:447-449 (1996).
- Vidalvanaclocha F, Alvarez A, Asumendi A, Urcelay B, Tonino P, Dinarello CA: Interleukin 1 (IL-1)-dependent melanoma hepatic metastasis *in vivo*; Increased endothelial adherence by IL-1-induced mannose receptors and growth factor production *in vitro*. *J Nat Cancer Inst* 88:198-205 (1996).
- Vijaykumar V, Topp EM: Diffusion of an anti-transferrin receptor antibody in cultured murine melanoma cell layers. *Pharmaceut Res* 12:1907-1916 (1995).
- Vondran T, Apel KH, Schmitz H: The infrared receptor of *Melanophila acuminata* De Geer (Coleoptera: Buprestidae): Ultrastructural study of a unique insect thermoreceptor and its possible descent from a hair mechanoreceptor. *Tissue Cell* 27:645-658 (1995).
- Waterhouse RN, Collier TL, O'Brien JC: Synthesis of 1-(trans-[I-123]iodopropen-2-yl)-4-(4-cyanophenoxymethyl)-piperidine: A selective sigma receptor radioligand for SPECT. *J Label Compound Radiopharm* 38:215-226 (1996).
- Wintzen M, Gilchrest BA: Proopiomelanocortin, its derived peptides, and the skin. *J Invest Dermatol* 106:3-10 (1996).
- Yang RB, Fulle HJ, Garbers DL: Chromosomal localization and genomic organization of genes encoding guanylyl cyclase receptors expressed in olfactory sensory neurons and retina. *Genomics* 31:367-372 (1996).
- Yannelli JR, Hyatt C, Mcconnell S, Hines K, Jacknin L, Parker L, Sanders M, Rosenberg SA: Growth of tumor-infiltrating lymphocytes from human solid cancers: Summary of a 5-year experience. *Int J Cancer* 65:413-421 (1996).

TYROSINASE, TYROSINASE RELATED PROTEINS & MOLECULAR BIOLOGY

- Argeson AC, Nelson KK, Siracusa LD: Molecular basis of the pleiotropic phenotype of mice carrying the hypervariable yellow (A^{hy}) mutation at the agouti locus. *Genetics* 142:557-567 (1996).
- Bailin T, Lee ST, Spritz RA: Genomic organization and sequence of D12S53E (Pmel 17), the human homologue of the mouse *silver* (*si*) locus. *J Invest Dermatol* 106:24-27 (1996).
- Bragadottir R, Jarkman S: Effects of PhXA41, a prostaglandin analogue, and PGF(2 α), on the corneal and intraretinal dc electroretinogram (ERG) of the albino rabbit eye. *Curr Eye Res* 14:1073-1080 (1995).
- Breimer LH, Winder AF, Panayiotidis P, Jay M, Moore A, Jay B: A trinucleotide deletion together with a base duplication event at codon 439 in the human tyrosinase gene identifies a mutational hotspot. *Clin Chim Acta* 243:35-42 (1995).
- Brichard VG, Herman J, Vanpel A, Wildmann C, Gaugler B, Wolfel T, Boon T, Lethe B: A tyrosinase nonapeptide presented by HLA-B44 is recognized on a human melanoma by autologous cytolytic T lymphocytes. *Eur J Immunol* 26:224-230 (1996).
- Eberle J, Garbe C, Wang NP, Orfanos CE: Incomplete expression of the tyrosinase gene family (Tyrosinase, TRP-1, and TRP-2) in human malignant melanoma cells *in vitro*. *Pigm Cell Res* 8:307-313 (1995).
- Ferguson CA, Kidson SH: Characteristic sequences in the promoter region of the chicken tyrosinase-encoding gene. *Gene* 169:191-195 (1996).
- Funk RHW, Gehr J, Rohen JW: Short-term hemodynamic changes in episcleral arteriovenous anastomoses correlate with venous pressure and IOP changes in the albino rabbit. *Curr Eye Res* 15:87-93 (1996).
- Granholtm DE, Reese RN, Granholtm NH: Agouti alleles influence thiol concentrations in hair follicles and extrafollicular tissues of mice (A^y/a , A^{wj}/A^{wj} , a/a). *Pigm Cell Res* 8:302-306 (1995).

- Granholt DE, Reese RN, Granholt NH: Agouti alleles alter cysteine and glutathione concentrations in hair follicles and serum of mice (A^y/a , A^{wj}/A^{wj} , and a/a). *J Invest Dermatol* 106:559-563 (1996).
- Hiremath SV, Gouripur VV, Patil PA: The anti-inflammatory activity of some sulphonamides in albino rats. *Indian J Med Res* 103:120-125 (1996).
- Jiang JW, Sharma SD, Nakamura S, Lai JY, Fink JL, Hruby VJ, Hadley ME: The melanotropic peptide, $[Nle^4, D-Phe^7]\alpha$ -MSH, stimulates human melanoma tyrosinase activity and inhibits cell proliferation. *Pigm Cell Res* 8:314-323 (1995).
- Jones BH, Kim JH, Zemel MB, Woychik RP, Michaud EJ, Wilkison WO, Moustaid N: Upregulation of adipocyte metabolism by agouti protein: Possible paracrine actions in yellow mouse obesity. *Amer J Physiol-Endocrinol Met* 33:E192-E196 (1996).
- Kahn V: Effect of kojic acid on the oxidation of DL-DOPA, norepinephrine, and dopamine by mushroom tyrosinase. *Pigm Cell Res* 8:234-240 (1995).
- Kanda K, Sato T, Suzuki K, Ishii S, Ejiri S, Enei H: Relationships between tyrosinase activity and gill browning during preservation of *Lentinus edodes* fruit-bodies. *Biosci Biotechnol Biochem* 60:479-480 (1996).
- Keil M, Lungarella G, Cavarra E, Vaneven P, Martorana PA: A scanning electron microscopic investigation of genetic emphysema in tight-skin, pallid, and beige mice, three different C57 BL/6J mutants. *Lab Invest* 74:353-362 (1996).
- Kucera GT, Bortner DZ, Rosenberg MP: Overexpression of an agouti cDNA in the skin of transgenic mice recapitulates dominant coat color phenotypes of spontaneous mutants. *Dev Biol* 173:162-173 (1996).
- Lamoreux ML, Zhou BK, Rosembat S, Orlov SJ: The pink-eyed-dilution protein and the eumelanin pheomelanin switch: In support of a unifying hypothesis. *Pigm Cell Res* 8:263-270 (1995).
- Lee ZH, Hou L, Moellmann G, Kuklinska E, Antol K, Fraser M, Halaban R, Kwon BS: Characterization and subcellular localization of human Pmel 17/ silver, a 100-kDa (pre)melanosomal membrane protein associated with 5,6,-dihydroxyindole-2-carboxylic acid (DHICA) converting activity. *J Invest Dermatol* 106:605-610 (1996).
- Morpurgo M, Schiavon O, Caliceti P, Veronese FM: Covalent modification of mushroom tyrosinase with different amphiphilic polymers for pharmaceutical and biocatalysis applications. *Appl Biochem Biotechnol* 56:59-72 (1996).
- Motro B, Wojtowicz JM, Bernstein A, Vanderkooy D: Steel mutant mice are deficient in hippocampal learning but not long-term potentiation. *Proc Natl Acad Sci USA* 93:1808-1813 (1996).
- Nellaippan K, Sugumaran M: On the presence of prophenoloxidase in the hemolymph of the horseshoe crab, *Limulus*. *Comp Biochem Physiol [B]* 113:163-168 (1996).
- Qu TY, Dong K, Sugioka K, Yamadori T: Demonstration of direct input from the retina to the lateral habenular nucleus in the albino rat. *Brain Res* 709:251-258 (1996).
- Shucard DW, Specht CM: Fast habituation of the long-latency auditory evoked potential in the awake albino rat. *Evoked Potential* 100:78-84 (1996).
- Skipper JCA, Hendrickson RC, Gulden PH, Brichard V, Vanpel A, Chen Y, Shabanowitz J, Wolfel T, Slingluff CL, Boon T, Hunt DF, Engelhard VH: An HLA-A2-restricted tyrosinase antigen on melanoma cells results from posttranslational modification and suggests a novel pathway for processing of membrane proteins. *J Exp Med* 183:527-534 (1996).
- Takeuchi T, Tamate H, Saijoh Y: A pleiotropic gene which controls coat color and lethality in early development in the mouse. *Zool Sci* 12:675-681 (1995).
- Tief K, Schmidt A, Aguzzi A, Beermann F: Tyrosinase is a new marker for cell populations in the mouse neural tube. *Develop Dynam* 205:445-456 (1996).
- Tissir F, Champagne B, Klingalevan K, Levan G, Szpirer J, Szpirer C: Regional localization of the rat genes encoding the cAMP-specific phosphodiesterases 3 (Pde4d) and 4 (Pde4b) and the tyrosinase-related protein 1 (Tyrrp1). *Mamm Genome* 7:222-223 (1996).

MISCELLANEOUS

- Ahmed AKMF, Dong K, Setsu T, Yamadori T: Correlation between different types of retinal ganglion cells and their projection pattern in the albino rat. *Brain Res* 706:163-168 (1996).
- Bowler JM, Peart D, Purves RD, Carre DA, Macknight ADC, Civan MM: Electron probe X-ray microanalysis of rabbit ciliary epithelium. *Exp Eye Res* 62:131-139 (1996).
- Galliresta L, Ensigni M: An intrinsic time limit between genesis and death of individual neurons in the developing retinal ganglion cell layer. *J Neurosci* 16:2318-2324 (1996).
- Green ST, Wilson OF: The effect of hair color on the incorporation of methadone into hair in the rat. *J Anal Toxicol* 20:121-123 (1996).
- Hood DC, Cideciyan AV, Halevy DA, Jacobson SG: Sites of disease action in a retinal dystrophy with supernormal and delayed rod electroretinogram b-waves. *Vision Res* 36:889-901 (1996).
- Marcus RC, Wang LC, Mason CA: Retinal axon divergence in the optic chiasm: Midline cells are unaffected by the albino mutation. *Development* 122:859-868 (1996).

Nayar JK, Knight JW: Wounding increases intracellular encapsulation (melanization) of developing *Brugia malayi* (Nematoda: Filarioidea) larvae in thoracic muscles of *Anopheles quadrimaculatus*. *Comp Biochem Physiol [A]* 112:553-557 (1995).

Paquet I, Chouinard N, Rouabhia M: Cutaneous cell and extracellular matrix responses to ultraviolet-B irradiation. *J Cell Physiol* 166:296-304 (1996).

☞ Slominski A: Sunlight - Can it prevent as well as cause cancer? *Cancer Res* 56:925(1996).

Yang ZT, Zhao Z, Panjwani N: Gangliosides of migrating and nonmigrating corneal epithelium in organ and cell culture. *Invest Ophthalmol Visual Sci* 37:501-510 (1996).